arXiv:math/0405193v1 [math.OA] 11 May 2004

R-TRANSFORM THEORY ON A TOWER OF AMALGAMATED NONCOMMUTATIVE PROBABILITY SPACES

ILWOO CHO

Abstract. In [12], we observed Amalgamated R-transform Theory. Different from the original definition of Voiculescu and Speicher (in [10] and [1]), we define R-transforms of operator-valued random variable(s) by operator-valued formal series. By doing that we can establish Amalgamated R-transform calculus. In this paper, we will concentrate on observing R-transforms on a tower of amalgamated noncommutative probability spaces. We will consider the R-transform calculus and relation between R-transforms on the tower and distributions with respect to the tower. As application, we will observe the compatibility of the given tower and commuting ladders of amalgamated noncommutative probability spaces.

Voiculescu developed Free Probability Theory. Here, the classical concept of Independence in Probability theory is replaced by a noncommutative analogue called Freeness (See [8]). There are two approaches to study Free Probability Theory. One of them is the original analytic approach of Voiculescu (See [8] and [7]) and the other one is the combinatorial approach of Speicher and Nica (See [1], [6] and [7]).

To observe the free additive convolution and free multiplicative convolution of two distributions induced by free random variables in a noncommutative probability space (over $B = \mathbb{C}$), Voiculescu defined R-transforms and S-transforms, respectively. These show that to study distributions is to study certain (B-) formal series for arbitrary noncommutative indeterminants.

Speicher defined the free cumulants which are the main objects in the combinatorial approach of Free Probability Theory. And he developed free probability theory by using the combinatorics and lattice theory on collections of noncrossing partitions (See [7]). Also, Speicher considered the operator-valued free probability theory, which is also defined and observed analytically by Voiculescu, when $\mathbb{C}$ is replaced to an arbitrary algebra B (See [1] and [7]). Nica defined R-transforms of several random variables (See [6]). He defined these R-transforms as multivariable formal series in noncommutative several indeterminants. To observe the R-transform, the Möbius Inversion under the embedding of lattices plays a key role (See [1],[7],[5],[9],[13] and [17]).

In [9], [19] and [20], we observed the amalgamated R-transform calculus. Actually, amalgamated R-transforms are defined originally by Voiculescu (See [7]) and are characterized combinatorially by Speicher (See [1]). In [9], we defined amalgamated R-transforms slightly different, compared with those in [1] and [7], because

---

Key words and phrases. Noncommutative Probability Spaces with Amalgamation, R-transforms, Moment Series, Operator-Valued Distributions, Towers of Algebras, Towers of amalgamated noncommutative probability spaces.

of the bimodule map property. We defined them as B-formal series, in Θ_B^s, in s-noncommutative indeterminents and tried to characterize, like in [6] and [7]. The main tool which is considered, for studying amalgamated R-transform calculus, is the amalgamated boxed convolution, $\boxed{*}_B$ defined in [9]. After defining boxed convolution over an arbitrary algebra B, we could get that

$$R_{x_1,...,x_s} \boxed{*}_B R_{y_1,...,y_s}^{symm(1_B)} = R_{x_1y_1,...,x_sy_s}, \text{ for any } s \in \mathbb{N},$$

where x_j's and y_j's are free B-valued random variables. However, different from scalar-valued case (in [6] and [7]), in general,

$$R_{x_1,...,x_s} \ \boxed{*}_B \ R_{y_1,...,y_s} \neq R_{x_1y_1,...,x_sy_s},$$

even if x_j's and y_j's are free over B.

In this paper, we will consider a tower of algebras. i.e

$$\mathbb{C} \subset B \subset A_1 \subset A_2 \subset .... \subset A_\infty,$$

where A_1 is an algebra over B and A_{j+1} is an algebra over A_j, for all $j = 1, 2, \cdots$, and A_∞ is an enveloping algebra of this tower of algebra. Notice that A_n is an algebra over B, again, for all $n \in \mathbb{N}$.

In Chapter 1, we will observe Amalgamated R-transform Theory established in [9]. In Chapter 2, we will consider the R-transform theory on a tower of algebras. In Chapter 3, we will study the relation between R-transforms on a tower of algebras and operator-valued (or scalar-valued) distributions on that tower. In Chapter 4, as application, we will define a operator-valued semicircularity on a tower of algebras and observe their R-transform theory. Also, after assuming the compatibility on the tower, we will observe the R-transform calculus and operator-valued distributions.

1. PRELIMINARIES

1.1. **Amalgamated Free Probability Theory.**

In this section, we will summarize and introduced the basic results from [1] and [9]. Throughout this section, let B be a unital algebra. The algebraic pair (A, φ) is said to be a noncommutative probability space over B (shortly, NCPSpace over B) if A is an algebra over B (i.e $1_B = 1_A \in B \subset A$) and $\varphi : A \to B$ is a B-functional (or a conditional expectation) ; φ satisfies

$$\varphi(b) = b, \text{ for all } b \in B$$

and

$$\varphi(bxb') = b\varphi(x)b', \text{ for all } b, b' \in B \text{ and } x \in A.$$

Let (A, φ) be a NCPSpace over B. Then, for the given B-functional, we can determine a moment multiplicative function $\widehat{\varphi} = (\varphi^{(n)})_{n=1}^{\infty} \in I(A, B)$, where

$$\varphi^{(n)}(a_1 \otimes ... \otimes a_n) = \varphi(a_1....a_n),$$

for all $a_1 \otimes ... \otimes a_n \in A^{\otimes_B n}$, $\forall n \in \mathbb{N}$.

We will denote noncrossing partitions over $\{1, ..., n\}$ $(n \in \mathbb{N})$ by $NC(n)$. Define an ordering on $NC(n)$;

$\theta = \{V_1, ..., V_k\} \leq \pi = \{W_1, ..., W_l\} \overset{def}{\Leftrightarrow}$ For each block $V_j \in \theta$, there exists only one block $W_p \in \pi$ such that $V_j \subset W_p$, for $j = 1, ..., k$ and $p = 1, ..., l$.

Then $(NC(n), \leq)$ is a complete lattice with its minimal element $0_n = \{(1), ..., (n)\}$ and its maximal element $1_n = \{(1, ..., n)\}$. We define the incidence algebra I_2 by a set of all complex-valued functions η on $\cup_{n=1}^{\infty} (NC(n) \times NC(n))$ satisfying $\eta(\theta, \pi) = 0$, whenever $\theta \nleq \pi$. Then, under the convolution

$$* : I_2 \times I_2 \to \mathbb{C}$$

defined by

$$\eta_1 * \eta_2(\theta, \pi) = \sum_{\theta \leq \sigma \leq \pi} \eta_1(\theta, \sigma) \cdot \eta_2(\sigma, \pi),$$

I_2 is indeed an algebra of complex-valued functions. Denote zeta, Möbius and delta functions in the incidence algebra I_2 by ζ, μ and δ, respectively. i.e

$$\zeta(\theta, \pi) = \begin{cases} 1 & \theta \leq \pi \\ 0 & otherwise, \end{cases}$$

$$\delta(\theta, \pi) = \begin{cases} 1 & \theta = \pi \\ 0 & otherwise, \end{cases}$$

and μ is the $(*)$-inverse of ζ. Notice that δ is the $(*)$-identity of I_2. By using the same notation $(*)$, we can define a convolution between $I(A, B)$ and I_2 by

$$\widehat{f} * \eta\,(a_1, ..., a_n\,;\,\pi) = \sum_{\pi \in NC(n)} \widehat{f}(\pi)(a_1 \otimes ... \otimes a_n)\eta(\pi, 1_n),$$

where $\widehat{f} \in I(A, B)$, $\eta \in I_1$, $\pi \in NC(n)$ and $a_j \in A$ $(j = 1, ..., n)$, for all $n \in \mathbb{N}$. Notice that $\widehat{f} * \eta \in I(A, B)$, too. Let $\widehat{\varphi}$ be a moment multiplicative function in $I(A, B)$ which we determined before. Then we can naturally define a cumulant multiplicative function $\widehat{c} = (c^{(n)})_{n=1}^{\infty} \in I(A, B)$ by

$$\widehat{c} = \widehat{\varphi} * \mu \quad \text{or} \quad \widehat{\varphi} = \widehat{c} * \zeta.$$

This says that if we have a moment multiplicative function, then we always get a cumulant multiplicative function and vice versa, by $(*)$. This relation is so-called "Möbius Inversion". More precisely, we have

$$\begin{aligned}\varphi(a_1...a_n) &= \varphi^{(n)}(a_1 \otimes ... \otimes a_n) \\ &= \sum_{\pi \in NC(n)} \widehat{c}(\pi)(a_1 \otimes ... \otimes a_n)\zeta(\pi, 1_n) \\ &= \sum_{\pi \in NC(n)} \widehat{c}(\pi)(a_1 \otimes ... \otimes a_n),\end{aligned}$$

for all $a_j \in A$ and $n \in \mathbb{N}$. Or equivalently,

$$c^{(n)}(a_1 \otimes ... \otimes a_n) \quad = \sum_{\pi \in NC(n)} \widehat{\varphi}(\pi)(a_1 \otimes ... \otimes a_n)\mu(\pi, 1_n).$$

Now, let (A_i, φ_i) be NCPSpaces over B, for all $i \in I$. Then we can define a amalgamated free product of A_i 's and amalgamated free product of φ_i's by

$$A \equiv *_B A_i \quad \text{and} \quad \varphi \equiv *_i \varphi_i,$$

respectively. Then, by Voiculescu, (A, φ) is again a NCPSpace over B and, as a vector space, A can be represented by

$$A = B \oplus \left(\oplus_{n=1}^{\infty} \left(\underset{i_1 \neq ... \neq i_n}{\oplus} (A_{i_1} \ominus B) \otimes ... \otimes (A_{i_n} \ominus B) \right) \right),$$

where $A_{i_j} \ominus B = \ker \varphi_{i_j}$. We will use Speicher's combinatorial definition of amalgamated free product of B-functionals ;

Definition 1.1. *Let (A_i, φ_i) be NCPSpaces over B, for all $i \in I$. Then $\varphi = *_i \varphi_i$ is the amalgamated free product of B-functionals φ_i's on $A = *_B A_i$ if the cumulant multiplicative function $\widehat{c} = \widehat{\varphi} * \mu \in I(A, B)$ has its restriction to $\underset{i \in I}{\cup} A_i$, $\underset{i \in I}{\oplus} \widehat{c_i}$, where $\widehat{c_i}$ is the cumulant multiplicative function induced by φ_i, for all $i \in I$ and, for each $n \in \mathbb{N}$,*

$$c^{(n)}(a_1 \otimes ... \otimes a_n) = \begin{cases} c_i^{(n)}(a_1 \otimes ... \otimes a_n) & \text{if } \forall a_j \in A_i \\ 0_B & \text{otherwise.} \end{cases}$$

Now, we will observe the freeness over B ;

Definition 1.2. *Let (A, φ)be a NCPSpace over B.*

(1) Subalgebras containing B, $A_i \subset A$ $(i \in I)$ are free (over B) if we let $\varphi_i = \varphi \mid_{A_i}$, for all $i \in I$, then $*_i \varphi_i$ has its cumulant multiplicative function $\widehat{c}$ such that its restriction to $\underset{i \in I}{\cup} A_i$ is $\underset{i \in I}{\oplus} \widehat{c_i}$, where $\widehat{c_i}$ is the cumulant multiplicative function induced by each φ_i, for all $i \in I$.

(2) Sebsets X_i $(i \in I)$ are free (over B) if subalgebras A_i's generated by B and X_i's are free in the sense of (1). i.e If we let $A_i = A\lg\left(X_i, B\right)$, for all $i \in I$, then A_i's are free over B.

In [1], Speicher showed that the above combinatorial freeness with amalgamation can be used alternatively with respect to Voiculescu's original freeness with amalgamation.

Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s$ be B-valued random variables $(s \in \mathbb{N})$. Define $(i_1, ..., i_n)$-th moment of $x_1, ..., x_s$ by

$$\varphi(x_{i_1} b_{i_2} x_{i_2} ... b_{i_n} x_{i_n}),$$

for arbitrary $b_{i_2}, ..., b_{i_n} \in B$, where $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$. Similarly, define a symmetric $(i_1, ..., i_n)$-th moment by the fixed $b_0 \in B$ by

$$\varphi(x_{i_1} b_0 x_{i_2} ... b_0 x_{i_n}).$$

If $b_0 = 1_B$, then we call this symmetric moments, trivial moments.

Cumulants defined below are main tool of combinatorial free probability theory ; in [9], we defined the $(i_1, ..., i_n)$-th cumulant of $x_1, ..., x_s$ by

$$k_n(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_{i_2} x_{i_2} \otimes ... \otimes b_{i_n} x_{i_n}),$$

for $b_{i_2}, ..., b_{i_n} \in B$, arbitrary, and $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$, where $\widehat{c} = (c^{(n)})_{n=1}^{\infty}$ is the cumulant multiplicative function induced by φ. Notice that, by Möbius inversion, we can always take such B-value whenever we have $(i_1, ..., i_n)$-th moment of $x_1, ..., x_s$. And, vice versa, if we have cumulants, then we can always take moments. Hence we can define a symmetric $(i_1, ..., i_n)$-th cumulant by $b_0 \in B$ of $x_1, ..., x_s$ by

$$k_n^{symm(b_0)}(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_0 x_{i_2} \otimes ... \otimes b_0 x_{i_n}).$$

If $b_0 = 1_B$, then it is said to be trivial cumulants of $x_1, ..., x_s$.

By Speicher, it is shown that subalgebras A_i $(i \in I)$ are free over B if and only if all mixed cumulants vanish.

Proposition 1.1. *(See [1] and [9]) Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s \in (A, \varphi)$ be B-valued random variables $(s \in \mathbb{N})$. Then $x_1, ..., x_s$ are free if and only if all their mixed cumulants vanish.* $\square$

1.2. **Amalgamated R-transform Theory.**

In this section, we will define an R-transform of several B-valued random variables. Note that to study R-transforms is to study operator-valued distributions. R-transforms with single variable is defined by Voiculescu (over B, in particular, $B = \mathbb{C}$. See [8] and [7]). Over $\mathbb{C}$, Nica defined multi-variable R-transforms in [6]. In [9], we extended his concepts, over B. R-transforms of B-valued random variables can be defined as B-formal series with its $(i_1, ..., i_n)$-th coefficients, $(i_1, ..., i_n)$-th cumulants of B-valued random variables, where $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$.

Definition 1.3. *Let (A, φ) be a NCPSpace over B and let $x_1, ..., x_s \in (A, \varphi)$ be B-valued random variables ($s \in \mathbb{N}$). Let $z_1, ..., z_s$ be noncommutative indeterminants. Define a moment series of $x_1, ..., x_s$, as a B-formal series, by*

$$M_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,..,i_n \in \{1,...,s\}} \varphi(x_{i_1} b_{i_2} x_{i_2} ... b_{i_n} x_{i_n})\, z_{i_1} ... z_{i_n},$$

where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary for all $(i_2, ..., i_n) \in \{1, ..., s\}^{n-1}$, $\forall n \in \mathbb{N}$.

Define an R-transform of $x_1, ..., x_s$, as a B-formal series, by

$$R_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} k_n(x_{i_1}, ..., x_{i_n})\, z_{i_1} ... z_{i_n},$$

with

$$k_n(x_{i_1}, ..., x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_{i_2} x_{i_2} \otimes ... \otimes b_{i_n} x_{i_n}),$$

where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary for all $(i_2, ..., i_n) \in \{1, ..., s\}^{n-1}$, $\forall n \in \mathbb{N}$. *Here, $\widehat{c} = (c^{(n)})_{n=1}^{\infty}$ is a cumulant multiplicative function induced by φ in $I(A, B)$.*

Denote a set of all B-formal series with s-noncommutative indeterminants ($s \in \mathbb{N}$), by Θ_B^s. i.e if $g \in \Theta_B^s$, then

$$g(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} b_{i_1,...,i_n}\, z_{i_1} ... z_{i_n},$$

where $b_{i_1,...,i_n} \in B$, for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $\forall n \in \mathbb{N}$. Trivially, by definition, $M_{x_1,...,x_s}, R_{x_1,...,x_s} \in \Theta_B^s$. By $\mathcal{R}_B^s$, we denote a set of all R-transforms of s-B-valued random variables. Recall that, set-theoratically,

$$\Theta_B^s = \mathcal{R}_B^s, \text{ sor all } s \in \mathbb{N}.$$

We can also define symmetric moment series and symmetric R-transform by $b_0 \in B$, by

$$M_{x_1,...,x_s}^{symm(b_0)}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} \varphi(x_{i_1} b_0 x_{i_2} ... b_0 x_{i_n})\, z_{i_1} ... z_{i_n}$$

and

$$R^{symm(b_0)}_{x_1,...,x_s}(z_1,...,z_s) = \sum_{n=1}^{\infty} \sum_{i_1,..,i_n \in \{1,...,s\}} k^{symm(b_0)}_{n}(x_{i_1},...,x_{i_n})\, z_{i_1}...z_{i_n},$$

with

$$k^{symm(b_0)}_{n}(x_{i_1},...,x_{i_n}) = c^{(n)}(x_{i_1} \otimes b_0 x_{i_2} \otimes ... \otimes b_0 x_{i_n}),$$

for all $(i_1,...,i_n) \in \{1,...,s\}^n$, $\forall n \in \mathbb{N}$.

If $b_0 = 1_B$, then we have trivial moment series and trivial R-transform of $x_1,...,x_s$ denoted by $M^t_{x_1,...,x_s}$ and $R^t_{x_1,...,x_s}$, respectively.

The followings are known in [1] and [9] ;

Proposition 1.2. *Let (A, φ) be a NCPSpace over B and let $x_1,...,x_s, y_1,...,y_p \in (A, \varphi)$ be B-valued random variables, where $s, p \in \mathbb{N}$. Suppose that $\{x_1,...,x_s\}$ and $\{y_1,...,y_p\}$ are free in (A, φ). Then*

(1) $R_{x_1,...,x_s,y_1,...,y_p}(z_1,...,z_{s+p}) = R_{x_1,...,x_s}(z_1,...,z_s) + R_{y_1,...,y_p}(z_1,...,z_p)$.

(2) If $s = p$, then $R_{x_1+y_1,...,x_s+y_s}(z_1,...,z_s) = (R_{x_1,...,x_s} + R_{y_1,...,y_s})(z_1,...,z_s)$.
$\square$

The above proposition is proved by the characterization of freeness with respect to cumulants. i.e $\{x_1,...,x_s\}$ and $\{y_1,...,y_p\}$ are free in (A, φ) if and only if their mixed cumulants vanish. Thus we have

$$\begin{aligned} k_n(p_{i_1},...,p_{i_n}) &= c^{(n)}(p_{i_1} \otimes b_{i_2} p_{i_2} \otimes ... \otimes b_{i_n} p_{i_n}) \\ &= (\widehat{c_x} \oplus \widehat{c_y})^{(n)}\,(p_{i_1} \otimes b_{i_2} p_{i_2} \otimes ... \otimes b_{i_n} p_{i_n}) \\ &= \begin{cases} k_n(x_{i_1},...,x_{i_n}) & or \\ k_n(y_{i_1},...,y_{i_n}) \end{cases} \end{aligned}$$

and if $s = p$, then

$$\begin{aligned} & k_n(x_{i_1} + y_{i_1},...,x_{i_n} + y_{i_n}) \\ &= c^{(n)}\left((x_{i_1} + y_{i_1}) \otimes b_{i_2}(x_{i_2} + y_{i_2}) \otimes ... \otimes b_{i_n}(x_{i_n} + y_{i_n})\right) \\ &= c^{(n)}(x_{i_1} \otimes b_{i_2} x_{i_2} \otimes ... \otimes b_{i_n} x_{i_n}) + c^{(n)}(y_{i_1} \otimes b_{i_2} y_{i_2} \otimes ... \otimes b_{i_n} y_{i_n}) + [Mixed] \end{aligned}$$

where $[Mixed]$ is the sum of mixed cumulants of x_j's and y_i's, by the bimodule map property of $c^{(n)}$

$$= k_n(x_{i_1},...,x_{i_n}) + k_n(y_{i_1},...,y_{i_n}) + 0_B.$$

Note that if $f, g \in \Theta^s_B$, then we can always choose free $\{x_1,...,x_s\}$ and $\{y_1,...,y_s\}$ in (some) NCPSpace over B, (A, φ), such that

$$f = R_{x_1,...,x_s} \quad \text{and} \quad g = R_{y_1,...,y_s}.$$

Definition 1.4. *(1) Let $s \in \mathbb{N}$. Let $(f, g) \in \Theta^s_B \times \Theta^s_B$. Define $\boxed{*} : \Theta^s_B \times \Theta^s_B \to \Theta^s_B$ by*

$$(f,g)=(R_{x_1,\ldots,x_s},\,R_{y_1,\ldots,y_s})\longmapsto R_{x_1,\ldots,x_s}\,\boxed{*}\,R_{y_1,\ldots,y_s}.$$

Here, $\{x_1,...,x_s\}$ *and* $\{y_1,...,y_s\}$ *are free in* (A,φ)*. Suppose that*

$$coef_{i_1,..,i_n}\left(R_{x_1,\ldots,x_s}\right)=c^{(n)}(x_{i_1}\otimes b_{i_2}x_{i_2}\otimes\ldots\otimes b_{i_n}x_{i_n})$$

and

$$coef_{i_1,\ldots,i_n}(R_{y_1,\ldots,y_s})=c^{(n)}(y_{i_1}\otimes b'_{i_2}y_{i_2}\otimes\ldots\otimes b'_{i_n}y_{i_n}),$$

for all $(i_1,...,i_n)\in\{1,...,s\}^n$, $n\in\mathbb{N}$, *where* $b_{i_j}, b'_{i_n}\in B$ *arbitrary. Then*

$$coef_{i_1,\ldots,i_n}\left(R_{x_1,\ldots,x_s}\,\boxed{*}\,R_{y_1,\ldots,y_s}\right)$$

$$=\sum_{\pi\in NC(n)}\left(\widehat{c_x}\oplus\widehat{c_y}\right)\left(\pi\cup Kr(\pi)\right)(x_{i_1}\otimes y_{i_1}\otimes b_{i_2}x_{i_2}\otimes b'_{i_2}y_{i_2}\otimes\ldots\otimes b_{i_n}x_{i_n}\otimes b'_{i_n}y_{i_n})$$

$$\overset{denote}{=}\sum_{\pi\in NC(n)}\left(k_\pi\oplus k_{Kr(\pi)}\right)(x_{i_1},y_{i_1},...,x_{i_n}y_{i_n}),$$

where $\widehat{c_x}\oplus\widehat{c_y}=\widehat{c}\mid_{A_x*_BA_y}$, $A_x=A\lg\left(\{x_i\}_{i=1}^s,B\right)$ *and* $A_y=A\lg\left(\{y_i\}_{i=1}^s,B\right)$ *and where* $\pi\cup Kr(\pi)$ *is an alternating union of partitions in* $NC(2n)$

Proposition 1.3. *(See [9])Let* (A,φ) *be a NCPSpace over* B *and let* $x_1,...,x_s,y_1,...,y_s\in(A,\varphi)$ *be* B*-valued random variables* ($s\in\mathbb{N}$)*. If* $\{x_1,...,x_s\}$ *and* $\{y_1,...,y_s\}$ *are free in* (A,φ)*, then we have*

$$k_n(x_{i_1}y_{i_1},...,x_{i_n}y_{i_n})$$

$$=\sum_{\pi\in NC(n)}\left(\widehat{c_x}\oplus\widehat{c_y}\right)\left(\pi\cup Kr(\pi)\right)(x_{i_1}\otimes y_{i_1}\otimes b_{i_2}x_{i_2}\otimes y_{i_2}\otimes\ldots\otimes b_{i_n}x_{i_n}\otimes y_{i_n})$$

$$\overset{denote}{=}\sum_{\pi\in NC(n)}\left(k_\pi\oplus k_{Kr(\pi)}^{symm(1_B)}\right)(x_{i_1},y_{i_1},...,x_{i_n},y_{i_n}),$$

for all $(i_1,...,i_n)\in\{1,...,s\}^n$, $\forall n\in\mathbb{N}$, $b_{i_2},...,b_{i_n}\in B$*, arbitrary, where* $\widehat{c_x}\oplus\widehat{c_y}=\widehat{c}\mid_{A_x*_BA_y}$, $A_x=A\lg\left(\{x_i\}_{i=1}^s,B\right)$ *and* $A_y=A\lg\left(\{y_i\}_{i=1}^s,B\right)$. $\square$

This shows that ;

Corollary 1.4. *(See [9]) Under the same condition with the previous proposition,*

$$R_{x_1,\ldots,x_s}\,\boxed{*}\,R^t_{y_1,\ldots,y_s}=R_{x_1y_1,\ldots,x_sy_s}.$$

$\square$

Notice that, in general, unless $b'_{i_2}=\ldots=b'_{i_n}=1_B$ in B,

$$R_{x_1,...,x_s} \boxed{*} R_{y_1,...,y_s} \neq R_{x_1y_1,...,x_sy_s}.$$

However, as we can see above,

$$R_{x_1,...,x_s} \boxed{*} R^t_{y_1,...,y_s} = R_{x_1y_1,...,x_sy_s}$$

and

$$R^t_{x_1,...,x_s} \boxed{*} R^t_{y_1,...,y_s} = R^t_{x_1y_1,...,x_sy_s},$$

where $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_s\}$ are free over B. Over $B = \mathbb{C}$, the last equation is proved by Nica and Speicher in [6] and [7]. Actually, their R-transforms (over $\mathbb{C}$) is our trivial R-transforms (over $\mathbb{C}$).

2. AMALGAMATED R-TRANSFORM THEORY ON A TOWER OF ALGEBRAS

2.1. **R-transform Calculus On a Tower of Amalgamated NCPSpaces.**

Let B be a unital algebra and let A_k's be algebras over B, for all $k = 1, 2, \cdots$. Moreover, assume that A_{j+1} is an algebra over A_j, for each $j = 1, 2, ...$. Then we can get a tower of algebras,

$$\mathbb{C} \subset B \subset A_1 \subset A_2 \subset \cdots.$$

Assume that there exists a sequence of conditional expectations $(\varphi_n : A_n \to A_{n-1})_{n=1}^{\infty}$, where $A_0 = B$. We can express this by

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} \cdots,$$

where $\varphi_0 : B \to \mathbb{C}$ is a fixed linear functional.

Definition 2.1. *Suppose that we have a tower of algebras,*

$$\mathbb{C} \subset B \subset A_1 \subset A_2 \subset A_3 \subset \cdots$$

such that A_j is an algebra over A_{j-1}, with $A_0 = B$, for each $j \in \mathbb{N}$. And assume that there is a sequence of conditional expectations, $\varphi_n : A_n \to A_{n-1}$, for all $n \in \mathbb{N}$. Then the tower of algebras together with conditional expectations,

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

is called a tower of amalgamated noncommutative probability spaces, where $\varphi_0 : B \to \mathbb{C}$ *is the fixed linear functional.*

If we fix $j \in \mathbb{N}$, then the inclusion $[A_j \subset^{\varphi_{j+1}} A_{j+1}]$ satisfies the R-transform theory observed in Chapter 1. i.e for each j, we can regard $[A_j \subset^{\varphi_{j+1}} A_{j+1}]$ as $[B \subset^{\varphi} A]$ observed in Chapter 1.

We are interested in the case $[\mathbb{C} \subset^{E_N} A_N]$, where $N \in \mathbb{N}$ and

$$E_N : A_N \to \mathbb{C}$$

is the linear functional defined by

$$E_N = \varphi_0 \circ \varphi_1 \circ \varphi_2 \circ \cdots \circ \varphi_N \overset{denote}{=} \varphi_0\varphi_1 ... \varphi_N.$$

Indeed, $E_N : A_N \to \mathbb{C}$ is a linear functional ;

$$\begin{aligned} E_N (\alpha x + \beta y) &= \varphi_0\varphi_1 ... \varphi_N (\alpha x + \beta y) \\ &= \varphi_0\varphi_1 ... \varphi_{N-1} (\alpha\varphi_N (x) + \beta\varphi_N(y)) \end{aligned}$$

by the bimodule map property of φ_N

$$= \varphi_0\varphi_1 ... \varphi_{N-2} \left(\alpha\varphi_{N-1}\varphi_N(x) + \beta\varphi_{N-1}\varphi_N(y)\right)$$

by the bimodule map property of φ_{N-1}

$$= ... = \alpha\varphi_0\varphi_1 ... \varphi_N(x) + \beta\varphi_0\varphi_1 ... \varphi_N(y)$$

by the linearity of φ_0.

So, $E_N = \varphi_0\varphi_1 ... \varphi_N$ is a well-determined linear functional on A_N, for each N.

Definition 2.2. *Let B be a unital algebra. Let $(\varphi_k : A_k \to A_{k-1})_{k=1}^{\infty}$, with $A_0 = B$ be a sequence of conditional expectations, where $\varphi_0 : B \to \mathbb{C}$ is a fixed linear functional. i.e, we have a tower of NCPSpaces,*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots.$$

For the fixed $j \in \mathbb{N}$, if $x_1, ..., x_s \in (A_{j+1}, \varphi_{j+1})$ are A_j-valued random variables ($s \in \mathbb{N}$), then the moment series of them is defined by

$$M^{(j+1)}_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} \varphi_{j+1} \left(x_{i_1} a_{i_2} x_{i_2} ... a_{i_n} x_{i_n}\right) z_{i_1} ... z_{i_n},$$

as a A_j-formal series in $\Theta^s_{A_j}$, where $b_{i_2}, ..., b_{i_b} \in A_j$ are arbitrary for each $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. Similarly, in $\Theta^s_{A_j}$, we can define a R-transform of $x_1, ..., x_s$ by

$$R^{(j+1)}_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} k_n^{(j+1)}\left(x_{i_1}, ..., x_{i_n}\right) z_{i_1}...z_{i_n},$$

with its $(i_1, ..., i_n)$-th coefficients

$$k_n^{(j+1)}\left(x_{i_1}, ..., x_{i_n}\right) = c^{(n)}_{j+1}\left(x_{i_1} \otimes a_{i_2}x_{i_2} \otimes ... \otimes a_{i_n}x_{i_n}\right),$$

where $a_{i_2}, ..., a_{i_n} \in A_j$ are arbitrary for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, and where $\widehat{c_{j+1}} = \left(c^{(k)}_{j+1}\right)_{k=1}^{\infty} \in I^c\left(A_{j+1}, A_j\right)$ is the cumulant multiplicative bimodule map induced by $\varphi_{j+1} : A_{j+1} \rightarrow A_j$. Similar to the $[B \subset^{\varphi} A]$-case in Chapter 1, we can define trivial moment series and trivial R-transform of $x_1, ..., x_s$ by

$$M^{(j+1)\,:\,t}_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} \varphi_{j+1}\left(x_{i_1}x_{i_2}...x_{i_n}\right) z_{i_1}...z_{i_n}$$

and

$$R^{(j+1)\,:\,t}_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} k_n^{(j+1)\,:\,t}\left(x_{i_1}, ..., x_{i_n}\right) z_{i_1}...z_{i_n},$$

in $\Theta^s_{A_j}$, as A_j-formal series, respectively. Let $E_{j+1} = \varphi_0\varphi_1 \cdots \varphi_{j+1} : A_{j+1} \rightarrow \mathbb{C}$ be a linear functional. Then a moment series of $x_1, ..., x_s$ and an R-transform of $x_1, ..., x_s$ are denoted by

$$m_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} E_{j+1}\left(x_{i_1}x_{i_2}...x_{i_n}\right) z_{i_1}...z_{i_n}$$

and

$$r_{x_1,...,x_s}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} k_n^{(E_{j+1})}\left(x_{i_1}, ..., x_{i_n}\right) z_{i_1}...z_{i_n},$$

as formal series in $\Theta^s_{\mathbb{C}} \overset{denote}{=} \Theta_s$, respectively. Here, the $(i_1, ..., i_n)$-th E_{j+1}-moments and $(i_1, ..., i_n)$-th E_{j+1}-cumulants are defined in the sense of Speicher and Nica (See [6] and [7]). i.e, by using our notation introduced in Chapter 1, they are just scalar-valued trivial moments and trivial cumulants.

Since we can regard E_{j+1} as a linear functional and $k_n^{(E_{j+1})}$ as a scalar-valued cumulants, we will use the notation used in [6] and [7], for the convenience. (For example, instead of using the notation for $\widehat{E_{j+1}}(\pi)(...)$ and $\widehat{c_{E_{j+1}}}(\pi)(...)$, we will use $E_{j+1\,:\,\pi}(...)$ and $k_\pi^{(E_{j+1})}(...)$, like in [6] and [7].)

Notice that we can apply all our results in Chapter 1 to $[A_j \subset^{\varphi_{j+1}} A_{j+1}]$, for the fixed $j \in \mathbb{N}$, in the tower of amalgamated NCPSpaces,

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots .$$

By $\boxed{*}_{A_j}$, we will denote the A_j-valued boxed convolution on $\Theta^s_{A_j}$, for all $s \in \mathbb{N}$. Also, we can consider the A_j-freeness for each j. In this section, we will concentrate on observing B-freeness of E_N and considering E_N-cumulants. But first, we can get the following R-transform calculus for each step $[A_j \subset^{\varphi_{j+1}} A_{j+1}]$. By definition and by Chapter 1, we have the following A_j-valued R-transform calculus, for each j-th step of a tower of NCPSpaces.

Proposition 2.1. *Let B be a unital algebra and assume that we have a tower of algebras $T_B\left((A_i)_{i=1}^{\infty}\right)$ and a sequence of conditional expectations $\left(\varphi_j : A_j \rightarrow A_{j-1}\right)_{j=1}^{\infty}$, with $A_0 = B$. i.e, we have a tower of NCPSpaces*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots .$$

Fix $j \in \mathbb{N}$. Let $x_1, ..., x_s, y_1, ..., y_s \in (A_{j+1}, \varphi_{j+1})$ be A_j-valued random variables ($s \in \mathbb{N}$). Assume that two subsets of A_{j+1}, $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ are free over A_j (in short, A_j-free). Then

(1) $R^{(j+1)}_{x_1,...,x_s,y_1,...,y_s}(z_1, ..., z_{2s}) = R^{(j+1)}_{x_1,...,x_s}(z_1, ..., z_s) + R^{(j+1)}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s}).$

(2) $R^{(j+1)}_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) = \left(R^{(j+!)}_{x_1,...,x_s} + R^{(j+1)}_{y_1,...,y_s}\right)(z_1, ..., z_s).$

(3) $R^{(j+1)}_{x_1y_1,...,x_sy_s}(z_1, ..., z_s) = \left(R^{(j+1)}_{x_1,...,x_s} \boxed{*}_{A_j} R^{(j+1)\,:\,t}_{y_1,...,y_s}\right)(z_1, ..., z_s).$ $\square$

From now, we will observe the E_{j+1}-cumulants of random variables. Notice that $(A_{j+1},\, E_{j+1})$ is a NCPSpace (over $\mathbb{C}$), since $E_{j+1} : A_{j+1} \rightarrow \mathbb{C}$ is a well-determined linear functional. Hence if $x_1, ..., x_s \in A_{j+1}$ are operators, then they can be regarded as (scalar-valued) random variables ($s \in \mathbb{N}$). So, E_{j+1}-cumulants of them are well-defined with respect to the multiplicative bimodule map $\widehat{c_{E_{j+1}}}$, induced by E_{j+1}.

Lemma 2.2. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Let $E_{j+1} = \varphi_0\varphi_1....\varphi_{j+1} : A_{j+1} \rightarrow B$ be a B-functional, for each $j = 1, ..., n$. Let $x_1, ..., x_s \in (A_{j+1}, E_{j+1})$ be B-valued random variables ($s \in \mathbb{N}$). Then

$k^{(E_{j+1})}_n\left(x_{i_1}, ..., x_{i_n}\right)$

$$= \sum_{\pi \in NC(n)} \left(\prod_{V=(v_1,...,v_k)\in\pi} \left(\sum_{\theta\in NC(k)} E_j(k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k})) \right) \right) \mu(\pi, 1_n),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$.

Proof. Fix $n \in \mathbb{N}$ and $(i_1, ..., i_n) \in \{1, ..., s\}^n$. Then

$$k_n^{(E_{j+1})}(x_{i_1}, ..., x_{i_n}) = c_{E_{j+1}}^{(n)}(x_{i_1} \otimes x_{i_2} \otimes ... \otimes x_{i_n})$$

by using of our notation (See Chapter 1)

$$= \sum_{\pi\in NC(n)} \widehat{E_{j+1}}(\pi)(x_{i_1} \otimes b_{i_2}x_{i_2} \otimes ... \otimes b_{i_n}x_{i_n})\ \mu(\pi, 1_n)$$

$$= \sum_{\pi\in NC(n)} E_{j+1\,:\,\pi}(x_{i_1}, ..., x_{i_n})\ \mu(\pi, 1_n)$$

by using notation of Nica and Speicher (See [6] and [7])

$$= \sum_{\pi\in NC(n)} \left(\prod_{V=(v_1,...,v_k)\in\pi} E_{j+1}(x_{v_1}...x_{v_k}) \right) \mu(\pi, 1_n).$$

(Since E_{j+1} is a linear functional, we don't need to consider the insertion property with respect to inner partitions and outer partitions. So, we can conclude the above last equality.)

Now, observe $E_{j+1}(x_{v_1}...x_{v_k})$, for the fixed block $V = (v_1, ..., v_k) \in \pi$, for $\pi \in NC(n)$;

$$\begin{aligned} E_{j+1}(x_{v_1}...x_{v_n}) &= \varphi_0\varphi_1...\varphi_j\varphi_{j+1}(x_{v_1}...x_{v_k}) \\ &= \varphi_0\varphi_1...\varphi_j\left(\varphi_{j+1}(x_{v_1}...x_{v_k})\right) \end{aligned}$$

(2.2.1)

$$= \varphi_0\varphi_1...\varphi_j \left(\sum_{\theta\in NC(k)} \widehat{c_{j+1}}(\theta)(x_{v_1} \otimes ... \otimes x_{v_k}) \right).$$

From now, we will denote a partition-dependent amalgamated **trivial** cumulant having the form, $\widehat{c_{j+1}}(\theta)(x_{v_1} \otimes ... \otimes x_{v_k})$, by $k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k})$. By using this new notation, we can re-express the formula (2.2.1) ;

(2.2.2)

$$= \varphi_0\varphi_1...\varphi_j \left(\sum_{\theta \in NC(k)} k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k}) \right)$$

$$= \varphi_0\varphi_1...\varphi_{j-1} \left(\sum_{\theta \in NC(k)} \varphi_j \left(k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k}) \right) \right)$$

$$= ....$$

$$= \sum_{\theta \in NC(k)} \varphi_0\varphi_1...\varphi_j \left(k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k}) \right)$$

$$= \sum_{\theta \in NC(k)} E_j \left(k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k}) \right).$$

Therefore,

$$k_n^{(E_{j+1})}(x_{i_1}, ..., x_{i_n}) = \sum_{\pi \in NC(n)} \left(\prod_{V=(v_1,...,v_k)\in\pi} E_{j+1}(x_{v_1}...x_{v_k}) \right) \mu(\pi, 1_n)$$

$$= \sum_{\pi \in NC(n)} \left(\prod_{V=(v_1,...,v_k)\in\pi} \left(\sum_{\theta \in NC(k)} E_j(k_{k\,:\,\theta}^{(j+1)\,:\,t}(x_{v_1}, ..., x_{v_k})) \right) \right) \mu(\pi, 1_n). \blacksquare$$

The above lemma shows the relation between E_{j+1}-cumulants and E_j-moments. Also, this shows that to compute the E_{j+1}-cumulants, we only need to consider the Möbius inversion for A_j-valued moments with respect to φ_{j+1}, for each block.

Corollary 2.3. *Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Define $E_{j+1} = \varphi_0\varphi_1...\varphi_j\varphi_{j+1}$, *for each* $j = 0, 1, ...$ *Let* $x_1, ..., x_s \in (A_{j+1}, E_{j+1})$ *be random variables* ($s \in \mathbb{N}$)*. Then*

$$k_n^{(E_{j+1})}(x_{i_1}, ..., x_{i_n}) = \sum_{\pi \in NC(n)} \left(\prod_{V=(v_1,...,v_k)\in\pi} E_j(\varphi_{j+1}(x_{v_1}...x_{v_k})) \right) \mu(\pi, 1_n),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. $\square$

Example 2.1. *Let* $j = 2$ *and let* $x, y \in (A_2, E_2)$ *be* B*-valued random variables. Then, by the straightforward computation, we have that*

$$k_3^{(E_2)}(x, x, y) = c^{(3)}(x \otimes x \otimes y)$$

$$= \sum_{\pi \in NC(3)} E_{2\,:\,\pi}\,(x, x, y)\,\mu(\pi, 1_3)$$

$$= E_2(xxy) - E_2(xx)E_2(y) - E_2\,(xy)\,E_2(x) \\ -E_2(x)E_2(xy) + E_2(x)E_2(x)E_2(y)$$

$$= \varphi_0\varphi_1\varphi_2\,(xxy) - \varphi_0\varphi_1\varphi_2(xx)\varphi_0\varphi_1\varphi_2(y) - \varphi_0\varphi_1\varphi_2\,(xy)\,\varphi_0\varphi_1\varphi_2(x) \\ -\varphi_0\varphi_1\varphi_2(x)\varphi_0\varphi_1\varphi_2(xy) + \varphi_0\varphi_1\varphi_2(x)\cdot\varphi_0\varphi_1\varphi_2(x)\cdot\varphi_0\varphi_1\varphi_2(y)$$

$$= E_1\left(\varphi_2(x^2y)\right) - E_1\left(\varphi_2(x^2)\right)E_1\left(\varphi_2(y)\right) - E_1\left(\varphi_2(xy)\right)E_1\left(\varphi_2(x)\right) \\ -E_1\left(\varphi_2(x)\right)E_1\left(\varphi_2(xy)\right) + E_1\left(\varphi_2(x)\right)E_1\left(\varphi_2(x)\right)E_1\left(\varphi_2(y)\right).$$

So, the above lemma and corollary are applied well in the above example. Here, by the above corollary, we have the informal way how to find such E_2*-cumulant ;*

(i) Consider the trivial A_1*-valued cumulant ;*

$$k_3^{(2)\,:\,t}\,(x, x, y) = \sum_{\pi \in NC(3)} \widehat{\varphi_2}(\pi)\,(x, x, y)\,\mu(\pi, 1_3)$$

$$= \varphi_2\left(x^2y\right) - \varphi_2(x)\varphi_2\,(xy) - \varphi_2\,(x\varphi_2(x)y) \\ -\varphi_2\left(x^2\right)\varphi_2(y) + \varphi_2(x)\varphi_2(x)\varphi_2(y).$$

(ii) Ignore the insertion property and act $E_1 = \varphi_0\varphi_1$*. Then we can get that*

$$k_3^{(E_2)}(x, x, y) = E_1\left(\varphi_2\left(x^2y\right)\right) - E_1\left(\varphi_2(x)\right)E_1\left(\varphi_2\,(xy)\right) \\ -E_1\left(\varphi_2\,(xy)\right)E_1\left(\varphi_2(x)\right) - E_1\left(\varphi_2(x^2)\right)E_1\left(\varphi_2(y)\right) \\ +E_1\left(\varphi_2(x)\right)E_1\left(\varphi_2(x)\right)E_1\left(\varphi_2(y)\right).$$

Corollary 2.4. *Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Let $E_{j+1} = \varphi_0\varphi_1...\varphi_{j+1}$ *be a linear functional on* A_{j+1}*, for all* j*. Let* $x \in (A_{j+1},\, E_{j+1})$ *be a random varialbe. Then*

$$k_n^{(E_{j+1})}\left(\underbrace{x, ..., x}_{n-times}\right) = \sum_{\pi \in NC(n)} \left(\prod_{V=(v_1,...,v_k)\in\pi} E_j\left(\varphi_{j+1}(x^k)\right)\right)\mu(\pi, 1_n).$$

□

Now, we will observe the R-transform calculus for E_{j+1}. The results are came from Chapter 1, [6] and [7] ;

Proposition 2.5. *(See [6] and [7]) Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Then, for any fixed $j \in \mathbb{N}$, *the following R-transform calculus holds true ; suppose that* $x_1, ..., x_s, y_1, ..., y_s \in (A_{j+1}, E_{j+1})$ *are random variables* ($s \in \mathbb{N}$) *and assume that* $X = \{x_1, ..., x_s\}$ *and* $Y = \{y_1, ..., y_s\}$ *are free. Then*

(1) $r_{x_1,...,x_s,y_1,...,y_s}(z_1, ..., z_{2s}) = r_{x_1,...,x_s}(z_1, ..., z_s) + r_{y_1,...,y_s}(z_{s+1}, ..., z_{2s}).$

(2) $r_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) = (r_{x_1,...,x_s} + r_{y_1,...,y_s})(z_1, ..., z_s).$ □

Proposition 2.6. *(See [6] and [7]) Let* B *be a unital algebra and let*

$$C \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Fix $j \in \mathbb{N}$. *Let* (A_{j+1}, E_{j+1}) *be a NCPSpace over* B, *with a linear functional,* $E_{j+1} = \varphi_0\varphi_1...\varphi_{j+1} : A_{j+1} \to \mathbb{C}$ *and let* $x_1, ..., x_s, y_1, ..., y_s \in (A_{j+1}, E_{j+1})$ *be* B*-valued random variables* ($s \in \mathbb{N}$). *If* $X = \{x_1, ..., x_s\}$ *and* $Y = \{y_1, ..., y_s\}$ *are free over* B, *then*

$$k_n^{(E_{j+1})}(x_{i_1}y_{i_1}, ..., x_{i_n}y_{i_n}) = \sum_{\pi \in NC(n)} \left(k_\pi^{(E_{j+1})}(x_{i_1}, ..., x_{i_n})\right)\left(k_{Kr(\pi)}^{(E_{j+1})}(y_{i_1}, ..., y_{i_n})\right),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. □

In the above proposition, $k_\pi^{(E_{j+1})}(...)$ and $k_{Kr(\pi)}^{(E_{j+1})}(...)$ are used as partition-dependent E_{j+1}-cumulants, in the sense of Speicher and Nica (See [6] and [7]). The above result can be proved by [6] and [7]. Also, it can be proved by using the result in Chapter 1.

Corollary 2.7. *(See [6] and [7]) Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Fix $j \in \mathbb{N}$. *Let* $X = \{x_1, ..., x_s\}$ *and* $Y = \{y_1, ..., y_s\}$ *be free subsets of random variables* ($s \in \mathbb{N}$) *in a NCPSpace,* (A_{j+1}, E_{j+1}). *Then*

$$r_{x_1y_1,...,x_sy_s}(z_1, ..., z_s) = (r_{x_1,...,x_s} \boxed{*}_s \, r_{y_1,...,y_s})(z_1, ..., z_s),$$

where $\boxed{*}_s : \Theta_s \times \Theta_s \to \Theta_s$ *is a (scalar-valued) boxed convolution in the sense of Nica and Speicher.* □

Define a conditional expectation

$$E_{k,\,j} : A_j \rightarrow A_{k-1},$$

by

$$E_{k,\,j} = \varphi_k \varphi_{k+1} ... \varphi_j \varphi_j,$$

for all $k < j$ in $\mathbb{N}$. Indeed, it is a conditional expectation, for the fixed $k < j$;

(i) Let $a \in A_{k-1}$. Then

$$E_{k,\,j}(a) = \varphi_k \varphi_{k+1} ... \varphi_{j-1} \left(\varphi_j(a) \right) = \varphi_k ... \varphi_{j-1}(a)$$

since $a \in A_{k-1} \subset A_{j-1}$

$$= ... = \varphi_k(a) = a.$$

So,

$$E_{k,\,j}(a) = a, \text{ for all } a \in A_{k-1}.$$

(ii) Let $a, a' \in A_{k-1}$ and $x \in A_j$. Then

$$\begin{aligned} E_{k,\,j}(axa') &= \varphi_k \varphi_{k+1} ... \varphi_{j-1} \left(\varphi_j(axa') \right) \\ &= \varphi_k \varphi_{k+1} ... \varphi_{j-1} \left(a \varphi_j(x) a' \right) \\ &= \varphi_k ... \varphi_{j-2} \left(\varphi_{j-1}(a \varphi_j(x) a') \right) \\ &= \varphi_k ... \varphi_{j-2} \left(a \cdot \varphi_{j-1} \varphi_j(x) \cdot a' \right) \\ &= ..... = a \cdot \varphi_k \varphi_{k+1} ... \varphi_j(x) \cdot a' = a E_{k,\,j}(x) a'. \end{aligned}$$

Thus we have that

$$E_{k,\,j}\left(axa'\right) = aE_{k,\,j}(x)a', \text{ for all } a, a' \in A_{k-1} \text{ and } x \in A_j.$$

By (i) and (ii), $E_{k,\,j} : A_j \rightarrow A_{k-1}$ is a conditional expectation (or a A_{k-1}-functional on A_j). And hence $(A_j,\ E_{k,\,j})$ is a NCPSpace over A_{k-1}. We can observe the following R-transform calculus, by Chapter 1 ; Defien, for A_{k-1}-valued random variables $x_1, ..., x_s$ in $(A_j,\, E_{k,\,j})$,

$$M_{x_1,\ldots,x_s}^{(E_{k,\,j})}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,\ldots,i_n \in \{1,\ldots,s\}} E_{k,\,j}\left(x_{i_1} a_{i_2} x_{i_2} ... a_{i_n} x_{i_n}\right) z_{i_1} ... z_{i_n}$$

and

$$R_{x_1,\ldots,x_s}^{(E_{k\,,\,j})}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,\ldots,i_n \in \{1,\ldots,s\}} k_n^{(E_{k,\,j})}\left(x_{i_1}, ..., x_{i_n}\right) z_{i_1} ... z_{i_n},$$

as the moment series of $x_1, ..., x_s$ and the R-transform of $x_1, ..., x_s$, respectively, in $\Theta_{A_{k-1}}^s$. Here, the $(i_1, ..., i_n)$-th cumulants of $x_1, .., x_s$ are determined by

$$k_n^{(E_{k,\,j})}\left(x_{i_1}, ..., x_{i_n}\right) = c_{E_{k,\,j}}^{(n)}\left(x_{i_1} \otimes a_{i_2}x_{i_2} \otimes ... \otimes a_{i_n}x_{i_n}\right),$$

where $a_{i_2}, ..., a_{i_n} \in A_{k-1}$ are arbitrary and $\widehat{c_{E_{k,\,j}}} = \left(c_{E_{k,\,j}}^{(m)}\right)_{m=1}^{\infty} \in I^c\left(A_j,\ A_{k-1}\right)$ is the cumulant multiplicative bimodule map induced by a A_{k-1}-functional, $E_{k,\,j}$. A trivial moment series of $x_1, ..., x_s$, denoted by $M_{x_1,...,x_s}^{(E_{k,\,j}):t}$, and a trivial R-transform of $x_1, ..., x_s$, denoted by $R_{x_1,...,x_s}^{(E_{k,\,j}):t}$, are defined usually.

Proposition 2.8. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset ...A_{k-1} \subset^{\varphi_k} A_k \subset ... \subset^{\varphi_j} A_j \subset^{\varphi_{j+1}} \cdots$$

be a tower of amalgamated NCPSpaces. Fix $k < j$ in $\mathbb{N}$. Define a conditional expectation $E_{k,\,j} : A_j \rightarrow A_{k-1}$, where $A_0 = B$. Let $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ be two subsets of A_{k-1}-valued random variables ($s \in \mathbb{N}$) in $\left(A_j,\ E_{k,\,j}\right)$. If X and Y are free over A_{k-1}, in $\left(A_j,\ E_{k,\,j}\right)$, then we have the following R-transform calculus ;

(1) $R_{x_1,...,x_s,y_1,...,y_s}^{(E_{k,\,j})}(z_1, ..., z_{2s}) = R_{x_1,...,x_s}^{(E_{k,\,j})}(z_1, ..., z_s) + R_{y_1,...,y_s}^{(E_{k,\,j})}(z_{s+1}, ..., z_{2s}).$

(2) $R_{x_1+y_1,...,x_s+y_s}^{(E_{k,\,j})}(z_1, ..., z_s) = \left(R_{x_1,...,x_s}^{(E_{k,\,j})} + R_{y_1,...,y_s}^{(E_{k,\,j})}\right)(z_1, ..., z_s).$

(3) $R_{x_1y_1,...,x_sy_s}^{(E_{k,\,j})}(z_1, ..., z_s) = \left(R_{x_1,...,x_s}^{(E_{k,\,j})} \boxed{\star}_{A_{k-1}}^{(E_{k,\,j})}\ R_{y_1,...,y_s}^{(E_{k,\,j})}\right)(z_1, ..., z_s),$

where $\boxed{\star}_{A_{k-1}} : \times_{A_{k-1}}^s \times \times_{A_{k-1}}^s \rightarrow \times_{A_{k-1}}^s$ is a A_{k-1}-valued boxed convolution on $\times_{A_{k-1}}^s$. $\square$

Clearly, $E_{k,\,k} = \varphi_k : A_k \rightarrow A_{k-1}$ is the A_{k-1}-functional.

Definition 2.3. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Define an enveloping algebra A_∞ and a linear functional $E_\infty : A_\infty \rightarrow \mathbb{C}$ by

$$E_\infty(x) = E_j(x), \ \ \textit{for } x \in A_j \subset A_\infty, \textit{ for all } \ j \in \mathbb{N}.$$

We will say the NCPSpace, (A_∞, E_∞), is the enveloping NCPSpace of the given tower.

By definition, for any $x \in A_N$ and $y \in A_M$ and for all $\alpha, \beta \in \mathbb{C}$, we have that

$$E_{\infty}\left(\alpha x+\beta y\right)=\alpha E_{\infty}(x)+\beta E_{\infty}(y)=\alpha E_{N}(x)+\beta E_{M}(y),$$

where $N, M \in \mathbb{N}$. Then above observation shows that there exits cumulants of random variables in (A_{∞}, E_{∞}) as follows ;

Definition 2.4. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces and (A_{∞}, E_{∞}) an enveloping NCPSpace of the tower. Let $x_1, ..., x_s \in (A_{\infty}, E_{\infty})$ be random variables ($s \in \mathbb{N}$). Define $(i_1, ..., i_n)$-th cumulants of $x_1, ..., x_s$ by

$$k_n^{(E_{\infty})}\left(x_{i_1}, ..., x_{i_n}\right)=\sum_{\pi \in NC(n)}\left(E_{\infty}\right)_{\pi}\left(x_{i_1}, ..., x_{i_n}\right)\mu(\pi, 1_n),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, where $k_n^{(E_{\infty})}(...)$ and $(E_{\infty})_{\pi}(...)$ are scalar-valued cumulants and partition-dependent moments in the sense of Speicher and Nica.

Suppose that $x, y \in (A_{\infty}, E_{\infty})$ are random variables and assume that $x \in A_j$ and $y \in A_k$, for some $j \leq k \in \mathbb{N}$. Then

$$\begin{aligned}
k_n^{(E_{\infty})}\left(x, y, x\right) &= \sum_{\pi \in NC(3)}\left(E_{\infty}\right)_{\pi}\left(x, y, x\right)\mu(\pi, 1_3) \\
&= E_{\infty}(xyx)-E_{\infty}(x)E_{\infty}(yx)-E_{\infty}(xy)E_{\infty}(x) \\
&\qquad -E_{\infty}\left(xE_{\infty}(y)x\right)+E_{\infty}(x)E_{\infty}(y)E_{\infty}(x) \\
&= E_{\infty}(xyx)-E_{\infty}(x)E_{\infty}(yx)-E_{\infty}(xy)E_{\infty}(x) \\
&\qquad -E_{\infty}\left(x^2\right)E_{\infty}(y)+E_{\infty}(x)E_{\infty}(y)E_{\infty}(x) \\
&= E_k(xyx)-E_j(x)E_k(yx)-E_k(xy)E_j(x) \\
&\qquad -E_j(x^2)E_k(y)+E_j(x)E_k(y)E_j(x).
\end{aligned}$$

Since $j \leq k$ and $A_j \subset^{E_{j,k}} A_k$, a random variable in (A_{∞}, E_{∞}),

$$xyx \in A_k, \;\; xy \in A_k \;\text{ and }\; yx \in A_k.$$

So, we can get the last equality, by definition of E_{∞}.

Proposition 2.9. *(See Chapter 1 or see [6] and [7]) Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces and (A_∞, E_∞), *an enveloping NCPSpace of this tower. Let* $X = \{x_1, ..., x_s\}$ *and* $Y = \{y_1, ..., y_s\}$ *be two free subsets of* (A_∞, E_∞), *then*

(2.9.1) $r^{(E_\infty)}_{x_1,...,x_s y_1,...,y_s}(z_1, ..., z_{2s}) = r^{(E_\infty)}_{x_1,...,x_s}(z_1, ..., z_s) + r^{(E_\infty)}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s}).$

(2.9.2) $r^{(E_\infty)}_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) = \left(r^{(E_\infty)}_{x_1,...,x_s} + r^{(E_\infty)}_{y_1,...,y_s}\right)(z_1, ..., z_s).$

(2.9.3) $r^{(E_\infty)}_{x_1 y_1,...,x_s y_s}(z_1, ..., z_s) = \left(r^{(E_\infty)}_{x_1,...,x_s} \boxed{*}\, r^{(E_\infty)}_{y_1,...,y_s}\right)(z_1, ..., z_s),$

where $\boxed{*} : \Theta_s \times \Theta_s \to \Theta_s$ *is the (scalar-valued) boxed convolution defined by Nica and Speicher.* $\square$

Now, we will observe the freeness on (A_∞, E_∞), more in detail.

Theorem 2.10. *Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces and (A_∞, E_∞), *an enveloping NCPSpace. Let* $X = \{x_1, ..., x_s\} \subset \cup_{j=1}^{k} A_j$ *and* $Y = \{y_1, ..., y_s\} \subset \cup_{j=k+1}^{l} A_j$ *be subsets of operators* ($s \in \mathbb{N}$), *where* $k < l$, *in* $\mathbb{N}$. *If* X *and* Y *are free in* $(A_l,\ E_l)$, *then (2.9.1), (2.9.2) and (2.9.3) holds true in* (A_∞, E_∞). *More precisely,*

(1) $r^{(E_\infty)}_{x_1,...,x_s,y_1,...,y_s}(z_1, ..., z_{2s}) = r^{(E_k)}_{x_1,...,x_s}(z_1, ..., z_s) + r^{(E_l)}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s}).$

(2) $r^{(E_\infty)}_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) = \left(r^{(E_k)}_{x_1,...,x_s} + r^{(E_l)}_{y_1,...,y_s}\right)(z_1, ..., z_s).$

(3) $r^{(E_\infty)}_{x_1 y_1,...,x_s y_s}(z_1, ..., z_s) = \left(r^{(E_k)}_{x_1,...,x_s} \boxed{*}_{\mathbb{C}}\ r^{(E_l)}_{y_1,...,y_s}\right)(z_1, ..., z_s).$

Proof. We can regards X and Y in (A_∞, E_∞) as subsets of random variables in (A_l, E_l). We have that

$$r^{(E_\infty)}_{x_1,...,x_s,y_1,...,y_s}(z_1, ..., z_{2s}) = r^{(E_l)}_{x_1,...,x_s,y_1...,y_s}(z_1, ..., z_{2s})$$

by definition of E_∞

$$= r^{(E_l)}_{x_1,...,x_s}(z_1, ..., z_s) + r^{(E_l)}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s})$$

by the freeness of X and Y, in (A_l, E_l)

$$= r^{(E_k)}_{x_1,...,x_s}(z_1, ..., z_s) + r^{(E_l)}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s})$$

since $E_l = \varphi_0\varphi_1 ... \varphi_k\varphi_{k+1} ... \varphi_l$ and $X \subset (A_k, E_k)$. Similarly, we have that

$$\begin{aligned} r^{(E_\infty)}_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) &= r^{(E_l)}_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) \\ &= \left(r^{(E_l)}_{x_1,...,x_s} + r^{(E_l)}_{y_1,...,y_s}\right)(z_1, ..., z_s) \\ &= \left(r^{(E_k)}_{x_1,...,x_s} + r^{(E_l)}_{y_1,...,y_s}\right)(z_1, ..., z_s). \end{aligned}$$

Now, fix $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. Notice that $x_1y_1, ..., x_sy_s$ can be regarded as random variables in (A_l, E_l). Consider the $(i_1, ..., i_n)$-th coefficient of $r^{(E_\infty)}_{x_1y_1,...,x_sy_s}$;

$$coef_{i_1,...,i_n}\left(r^{(E_\infty)}_{x_1y_1,...,x_sy_s}\right) = coef_{i_1,...,i_n}\left(r^{(E_l)}_{x_1y_1,...,x_sy_s}\right)$$

by the fact that $x_jy_j \in A_l$, for all j and by the definition of E_∞

$$\begin{aligned} &= k^{(E_l)}_n\left(x_{i_1}y_{i_1}, ..., x_{i_n}y_{i_n}\right) \\ &= \sum_{\pi \in NC(n)} k_\pi\left(x_{i_1}, ..., x_{i_n}\right) k_{Kr(\pi)}\left(y_{i_1}, ..., y_{i_n}\right) \end{aligned}$$

by the freeness of X and Y in (A_l, E_l), in the sense of Speicher and Nica (See Chapter 1 or see [6] and [7])

$$\begin{aligned} &= coef_{i_1,...,i_n}\left(r^{(E_l)}_{x_1,...,x_s} \boxed{*}\, r^{(E_l)}_{y_1,...,y_s}\right) \\ &= coef_{i_1,...,i_n}\left(r^{(E_k)}_{x_1,...,x_s} \boxed{*}\, r^{(E_l)}_{y_1,...,y_s}\right), \end{aligned}$$

since $x_1, ..., x_s \in \cup^k_{j=1} A_j$ and $E_l = E_k\varphi_{k+1} ... \varphi_l$. ∎

Corollary 2.11. *Let B be a unital algebra and suppose that we have a tower of amalgamated NCPSpaces, $\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} \cdots$. Let X and Y be subsets in A_k. If X and Y are free in (A_k, E_k), then X and Y are free in (A_N, E_N), for all $N > k$.* □

Remark 2.1. *Suppose that we choose $x_1, ..., x_s \in \cup^k_{j=l+1} A_j$ and $y_1, ..., y_s \in \cup^l_{j=1} A_j$, where $k > l$, in this time. Then we can get the same results as above, but it means different.*

Remark 2.2. *If we take a sufficiently large $N \in \mathbb{N}$, and if we replace ∞ to such N, then we can get the same results as in the previous theorem.*

More generally, we have that ;

Theorem 2.12. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Fix a sufficiently large $N \in \mathbb{N}$ and fix $k < N$. Consider the conditional expectation (or a A_{k-1}-functional), $E_{k,\,N} : A_N \to A_{k-1}$. Let $X = \{x_1, ..., x_s\} \subset \cup_{j=k}^{m} A_j$ and $Y = \{y_1, ..., y_s\} \subset \cup_{j=m+1}^{n} A_j$ be two subsets, in $(A_N,\, E_{k,N})$ of A_{k-1}-valued random variables ($s \in \mathbb{N}$), where $k < m < n < N$. If X and Y are free over A_{k-1}, in $(A_N, E_{k,N})$, then

(1) $R^{(E_{k,N})}_{x_1,...,x_s,y_1,...,y_s}(z_1, ..., z_{2s}) = R^{(E_{k,m})}_{x_1,...,x_s}(z_1, ..., z_s) + R^{(E_{k,n})}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s}).$

(2) $R^{(E_{k,N})}_{x_1+y_1,...,x_s+y_s}(z_1, ..., z_s) = \left(R^{(E_{k,m})}_{x_1,...,x_s} + R^{(E_{k,n})}_{y_1,...,y_s}\right)(z_1, ..., z_s).$

(3) $R^{(E_{k,N})}_{x_1y_1,...,x_sy_s}(z_1, ..., z_s) = \left(R^{(E_{k,m})}_{x_1,...,x_s} \boxed{*}_{A_{k-1}} R^{(E_{k,n})\,:\,t}_{y_1,...,y_s}\right)(z_1, ..., z_s).$

Proof. We can regard X and Y are subsets of A_N and consider them as A_{k-1}-valued random variables in $(A_N,\, E_{k,N})$. Since X and Y are free over A_{k-1}, we have that

$$R^{(E_{k,N})}_{x_1,...,x_s,y_1,...,y_s}(z_1, ..., z_{2s}) = R^{(E_{k,N})}_{x_1,...,x_s}(z_1, ..., z_s) + R^{(E_{k,N})}_{y_1,...,y_s}(z_{s+1}, ..., z_{2s})$$

Observe that

$$\begin{aligned} coef_{i_1,...,i_n}\left(R^{(E_{k,N})}_{x_1,...,x_s}\right) &= k_n^{(E_{k,N})}\left(x_{i_1}, ..., x_{i_n}\right) \\ &= c^{(n)}_{E_{k,N}}\left(x_{i_1} \otimes a_{i_2}x_{i_2} \otimes ... \otimes a_{i_n}x_{i_n}\right) \\ &= \sum_{\pi \in NC(n)} \widehat{E_{k,N}}(\pi)\left(x_{i_1} \otimes a_{i_2}x_{i_2} \otimes ... \otimes a_{i_n}x_{i_n}\right)\mu(\pi, 1_n) \\ &= \sum_{\pi \in NC(n)} \widehat{E_{k,m}}(\pi)\left(x_{i_1} \otimes a_{i_2}x_{i_2} \otimes ... \otimes a_{i_n}x_{i_n}\right)\,\mu(\pi, 1_n), \end{aligned}$$

since $E_{k,N} = E_{k,m}\varphi_{m+1}...\varphi_n...\varphi_N$ and x_{i_j}'s are in A_m, where $a_{i_2}, ..., a_{i_n} \in A_{k-1}$ are arbitrary. Thus

$$coef_{i_1,...,i_n}\left(R^{(E_{k,N})}_{x_1,...,x_s}\right) = coef_{i_1,...,i_n}\left(R^{(E_{k,m})}_{x_1,...,x_s}\right).$$

Similarly,

$$coef_{i_1,...,i_n}\left(R^{(E_{k,N})}_{y_1,...,y_s}\right) = coef_{i_1,...,i_n}\left(R^{(E_{k,n})}_{y_1,...,y_s}\right).$$

So, (1) is proved. By using the same idea, we can easily prove (2).

Now, fix $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. Then

$$\begin{aligned}
&coef_{i_1,...,i_p}\left(R^{(E_{k,N})}_{x_1y_1,...,x_sy_s}\right) = k_p^{(E_{k,N})}\left(x_{i_1}y_{i_1}, ..., x_{i_p}y_{i_p}\right) \\
&\quad = c^{(p)}_{E_{k,N}}\left(x_{i_1}y_{i_1} \otimes a_{i_2}x_{i_2}y_{i_2} \otimes ... \otimes a_{i_p}x_{i_p}y_{i_p}\right) \\
&\quad = \sum_{\pi \in NC(p)} \widehat{E_{k,N}}(\pi)\left(x_{i_1}y_{i_1} \otimes a_{i_2}x_{i_2}y_{i_2} \otimes ... \otimes a_{i_p}x_{i_p}y_{i_p}\right)\,\mu(\pi, 1_p) \\
&\quad = \sum_{\pi \in NC(p)} \left(\widehat{c_{E_{k,m}}} \oplus \widehat{c_{E_{k,n}}}\right)\left(x_{i_1} \otimes y_{i_1} \otimes a_{i_2}x_{i_2} \otimes y_{i_2} \otimes ... \otimes a_{i_p}x_{i_p} \otimes y_{i_p}\right)
\end{aligned}$$

by the A_{k-1}-freeness of X and Y in $(A_N,\, E_{k,N})$

$$= coef_{i_1,...,i_n}\left(R^{(E_{n,m})}_{x_1,...,x_s}\ \boxed{\ast}_{A_{k-1}}\ R^{(E_{k,n})\,:\,t}_{y_1,...,y_s}\right).$$

Therefore,

$$R^{(E_{k,N})}_{x_1y_1,...,x_sy_s} = R^{(E_{k,m})}_{x_1,...,x_s}\ \boxed{\ast}_B\ R^{(E_{k,n})}_{y_1,...,y_s}.$$

▮

Fix $k \in \mathbb{N}$. We can define an enveloping NCPSpace over A_{k-1}, $(A_\infty,\, E_{k,\infty})$, where A_∞ is an enveloping algebra of the tower and $E_{k,\infty} : A_\infty \to A_{k-1}$ is defined by

$$E_{k,\infty}(x) \overset{def}{=} E_{k,j}(x), \text{ for all } x \in A_j,$$

where $j > k$.

Corollary 2.13. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Let X and Y be a two subsets of A_k. If X and Y are free over A_{k-1}, in $(A_k, E_{k,k})$, then X and Y are free over A_{k-1}, in $(A_N, E_{k,N})$, for all $N > k$, in $\mathbb{N}$. □

Remark 2.3. *Let $(A_\infty,\, E_{k,\infty})$ be an enveloping NCPSpace over A_{k-1}. Then the same results with the previous theorem holds true when we replace N by ∞.*

2.2. **Distributions On a Tower of Algebras.**

In this section, we will consider (scalar-valued or operator-valued) distributions when the tower of amalgamated NCPSpaces is given. Throughout this section, we let

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Also, like in the previous section, we will use the following notations ;

$$E_N = \varphi_0\varphi_1...\varphi_N : A_N \to \mathbb{C}, \text{ a linear functional.}$$

For $k < N$,

$$E_{k,N} = \varphi_k\varphi_{k+1}...\varphi_N : A_N \to A_{k-1}, \text{ a } A_{k-1}\text{-functional}$$

with

$$E_{1,N} : A_N \to B, \text{ a } B\text{-functional.}$$

In the previous section, we considered the R-transform calculus of E_N and $E_{k,\,N}$. Also, when we consider the enveloping algebra of the given tower, A_∞, we could define a linear functional and A_{k-1}-functional, $E_\infty : A_\infty \to \mathbb{C}$ and $E_{k,\,\infty} : A_\infty \to A_{k-1}$, respectively. Also, R-transform calculus related to them are given in the previous section. Define

$$\sum\nolimits_s = \{\sigma : \mathbb{C}[X_1, ..., X_s] \to \mathbb{C} : \sigma \text{ is a } \mathbb{C}\text{-valued distribution}\}$$

and

$$\sum\nolimits_{A_{k-1}}^s = \{\sigma^{(A_{k-1})} : A_{k-1}[X_1, ..., X_s] \to A_{k-1} : \sigma^{(A_{k-1})} \text{ is a } A_{k-1}\text{-functional}\},$$

for the fixed $s \in \mathbb{N}$, where $X_1, ..., X_s$ are noncommutative indeterminents. If $\sigma \in \sum_s$, then there exists an arbitrary NCPSpace (A, φ) and random variables $x_1, ..., x_s \in (A, \varphi)$ such that

$$\sigma\,(P) = \varphi\,(P(x_1, ..., x_s)) \in \mathbb{C}, \text{ for all } P \in C[X_1, ..., X_s].$$

And if $\sigma^{(A_{k-1})} \in \sum_{A_{k-1}}^s$, then there exsits an arbitrary NCPSpace over A_{k-1}, (D, ψ), and A_{k-1}-valued random variables $y_1, ..., y_s \in (D, \psi)$ such that

$$\sigma^{(A_{k-1})}(Q) = \psi\,(Q(y_1, ..., y_s)) \in A_{k-1}, \text{ for all } Q \in A_{k-1}[X_1, ..., X_s].$$

In particular, we denote such σ and $\sigma^{(A_{k-1})}$ by

$$\sigma_{x_1,...,x_s \in (A,\varphi)} \quad \text{and} \quad \sigma^{(A_{k-1})}_{y_1,...,y_s \in (D,\psi)},$$

respectively. Notice that, for the given σ and $\sigma^{(A_{k-1})}$, $x_1, ..., x_s \in (A, \varphi)$ and $y_1, ..., y_s \in (D, \psi)$ are Not uniquely determined. So, we can regard them as equivalence class under the identically-distributedness. i.e

$$\begin{aligned} &\sigma = \sigma_{x_1,...,x_s \in (A,\varphi)} \sim \sigma_{x_1',...,x_s' \in (A',\varphi')} \\ &\qquad \Longleftrightarrow \varphi\left(P(x_1, ..., x_s)\right) = \varphi'\left(P(x_1', ..., x_s')\right), \ \forall \ P \in \mathbb{C}[X_1, ..., X_s] \end{aligned}$$

and

$$\begin{aligned} &\sigma^{(A_{k-1})} = \sigma^{(A_{k-1})}_{y_1,...,y_s \in (D,\psi)} \sim \sigma^{(A_{k-1})}_{y_1',...,y_s' \in (D',\psi')} \\ &\qquad \Longleftrightarrow \psi\left(Q(y_1, ..., y_s)\right) = \psi'\left(Q(y_1', ..., y_s')\right), \ \forall Q \in A_{k-1}[X_1, ..., X_s]. \end{aligned}$$

Recall the definitions ;

the free additive convolution $\boxplus : \sum_s \times \sum_s \rightarrow \sum_s$,

the free multiplicative convolution $\boxtimes : \sum_s \times \sum_s \rightarrow \sum_s$

and

the oprator-valued free additive convolution $\boxplus_{A_{k-1}} : \sum_{A_{k-1}}^s \times \sum_{A_{k-1}}^s \rightarrow \sum_{A_{k-1}}^s$.

(See [8] for $\boxplus$ and $\boxtimes$. See [1] for $\boxplus_{A_{k-1}}$) Now, we will define the oprerator-valued free multiplicative convolution of operator-valued distributions

$$\boxtimes_{A_{k-1}} : \sum\nolimits_{A_{k-1}}^s \times \sum\nolimits_{A_{k-1}}^s \rightarrow \sum\nolimits_{A_{k-1}}^s$$

by

$$\sigma^{(A_{k-1})}_{y_1,...,y_s \in (D,\psi)} \ \boxtimes_{A_{k-1}} \ \sigma^{(A_{k-1})}_{p_1,...,p_s \in (D',\psi')} = \sigma^{(A_{k-1})}_{y_1p_1,...,y_sp_s \in \left(D *_{A_{k-1}} D', \ \psi * \psi'\right)},$$

where $\left(D *_{A_{k-1}} D', \, \psi * \psi\right)$ is the free product of (D, ψ) and (D', ψ'), with amalgamation over A_{k-1}.

Rest of this section, we will consider the B-valued distributions depending on our tower.

Proposition 2.14. *Let* $\sigma \in \sum_s$ *and* $\sigma^{(A_{k-1})} \in \sum_{A_{k-1}}^s$. *Then*

(1) $\sigma_{x_1,...,x_s \in (A_\infty, E_\infty)} \ \boxplus \ \sigma_{y_1,...,y_s \in (A_\infty, E_\infty)} = \sigma_{x_1+y_1,...,x_s+y_s \in (A_\infty, E_\infty)}$.

(2) $\sigma^{(A_{k-1})}_{x_1,...,x_s \in (A_\infty, E_{k,\infty})} \ \boxplus_{A_{k-1}} \ \sigma^{(A_{k-1})}_{y_1,...,y_s \in (A_\infty, E_{k,\infty})} = \sigma^{(A_{k-1})}_{x_1+y_1,...,x_s+y_s \in (A_\infty, E_{k,\infty})}$.

(3) $\sigma_{x_1,...,x_s \in (A_j, E_j)} \ \boxplus \ \sigma_{y_1,...,y_s \in (A_k, E_k)} = \sigma_{x_1+y_1,...,x_s+y_s \in (A_k, E_k)}$, *if* $j < k$.

(4) $\sigma^{(A_{k-1})}_{x_1,...,x_s \in (A_i, \, E_{k,i})} \ \boxplus_{A_{k-1}} \sigma^{(A_{k-1})}_{y_1,...,y_s \in (A_j, \, E_{k,j})} = \sigma^{(A_{k-1})}_{x_1+y_1,...,x_s+y_s \in (A_j, E_{k,j})}$,

if $K < i < j$. $\square$

Proposition 2.15. *Let* $\sigma \in \sum_{s}$ *and* $\sigma^{(A_{k-1})} \in \sum_{A_{k-1}}$. *Then*

(1) $\sigma_{x_1,...,x_s \in (A_\infty, E_\infty)} \boxtimes \sigma_{y_1,...,y_s \in (A_\infty, E_\infty)} = \sigma_{x_1y_1,...,x_sy_s \in (A_\infty, E_\infty)}$.

(2) $\sigma^{(A_{k-1})}_{x_1,...,x_s \in (A_\infty, E_{k,\infty})} \boxtimes_{A_{k-1}} \sigma^{(A_{k-1})}_{y_1,...,y_s \in (A_\infty, E_{k,\infty})} = \sigma^{(A_{k-1})}_{x_1y_1,...,x_sy_s \in (A_\infty, E_{k,\infty})}$.

(3) $\sigma_{x_1,...,x_s \in (A_i, E_i)} \boxtimes \sigma_{y_1,...,y_s \in (A_j, E_j)} = \sigma_{x_1y_1,...,x_sy_s \in (A_j, E_j)}$, *if* $i < j$.

(4) $\sigma^{(A_{k-1})}_{x_1,...,x_s \in (A_i, E_{k,i})} \boxtimes_{A_{k-1}} \sigma^{(A_{k-1})}_{y_1,...,y_s \in (A_j, E_{k,j})} = \sigma^{(A_{k-1})}_{x_1y_1,...,x_sy_s \in (A_j,\, E_{k,j})}$,

if $k < i < j$.

Proof. Suppose that $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ are two subsets of (A_∞, E_∞) which are free. Then, there exists $i, j \in N$ such that

$$X \subset \cup_{l=1}^{m} A_l \subseteq A_i \quad \text{and} \quad Y \subset \cup_{l=1}^{n} A_l \subseteq A_j.$$

Assume that $m < n$. Then $\{x_1y_1, ..., x_sy_s\} \subset (A_j, E_j)$. By freeness,

$$\sigma_{x_1,...,x_s \in (A_i, E_i)} \boxtimes \sigma_{y_1,...,y_s \in (A_j, E_j)} = \sigma_{x_1y_1,...,x_sy_s \in (A_j, E_j)}.$$

So, (1) and (3) are proved. Similarly, $\{x_1y_1, ..., x_sy_s\} \subset (A_j, E_{k,j})$, if $k < i < j$. So,

$$\sigma^{(A_{k-1})}_{x_1,...,x_s \in (A_i, E_{k,i})} \boxtimes_{A_{k-1}} \sigma^{(A_{k-1})}_{y_1,...,y_s \in (A_j, E_{k,\,j})} = \sigma^{(A_{k-1})}_{x_1y_1,...,x_sy_s \in (A_j, E_{k,j})}.$$

Therefore, we can prove (2) and (4). ∎

Now, we will observe the following concept introduced in [18], so-called "compatibility". The R-transform theory, under this compatibility is studied in [12].

Definition 2.5. *Let* B *be a unital algebra and* A, *an algebra over* B. *Let* (A, φ) *be a NCPSpace and let* (A, E) *be a NCPSpace over* B, *with its* B-*functional* $E : A \to B$. *We say that* (A, φ) *and* (A, E) *are compatible if*

$$\varphi(x) = \varphi\left(E(x)\right), \quad \text{for all } x \in A.$$

As we can see in [12], even under the compatibility, to compute scalar-valued R-transforms from the operator-valued R-transforms or operator-valued moment series is complicated. But [12] provides the method.

Theorem 2.16. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. Then, for any fixed $k \in \mathbb{N}$, $(A_N,\, E_{k,N})$ and $(A_N,\, E_N)$ are compatible. In particular, if we have the enveloping algebra of the tower, A_∞, then $(A_\infty, E_{k,\infty})$ and (A_∞, E_∞) are compatible, for all k.

Proof. Fix $N \in \mathbb{N}$. By definition, we have that

$$\begin{aligned} E_N(x) &= \varphi_0\varphi_1 ... \varphi_N(x) \\ &= (\varphi_0\varphi_1 ... \varphi_{k-1})(\varphi_k ... \varphi_N)(x) \\ &= E_{k-1}\left(E_{k,N}(x)\right) \end{aligned}$$

and

$$\begin{aligned} E_N\left(E_{k,N}(x)\right) &= (\varphi_0\varphi_1 ... \varphi_N)\left(\varphi_k ... \varphi_N(x)\right) \\ &= \varphi_0\varphi_1 ... \varphi_N(x) = E_{k-1}\left(E_{k,N}(x)\right), \end{aligned}$$

for all $x \in A_N$. Therefore, for all x in A_N,

$$E_\infty(x) = E_N\left(E_{k,N}(x)\right).$$

∎

The above theorem shows that our tower of amalgamated NCPSpaces has nice compatible properties.

2.3. **Compressed R-transform Theory For the Chain of Projections.**

In this section, we will observe a tower of amalgamated NCPSpaces, generated by a chain of projections $(p_n)_{n=1}^{\infty}$ of a NCPSpace (A, φ). The B-valued R-transform theory of compressed algebras (i.e, $B \subset pAp$, where $p \in A$ is a projection) and B-valued freeness of them are studied in [10]. Also, by [35], there is the compatibility of $(A_k \subset A,\, E_{k+1,\,\infty})$ and $(A,\, E_k \circ E_{k+1,\,\infty})$. It is possible that there are only finitely many projections in the given chain (for example, an algebra A can be a finite dimensional algebra over a finite dimensional algebra B).

Definition 2.6. *Let A be a unital algebra. A nonzero element $p \in A \setminus \{0_A\}$ is called a projection (or an idempotent element) if p satisfies*

$$p^2 = p.$$

By A_{pro}*, we will denote a subset of all projections in an algebra,* A*. It is said that* $(p_n)_{n=1}^{\infty} \subset A_{pro}$ *is a chain of projections if*

$$\mathbb{C} \subset p_1 A p_1 \subset p_2 A p_2 \subset p_3 A p_3 \subset \cdots$$

is a tower of compressed subalgebras of A *satisfying the property that* $p_j A p_j$ *are subalgebras of* $p_{j+1} A p_{j+1}$*, for all* $j \in \mathbb{N}$*. For the convinience of using notation, we will denote maximal projection in a chain of projections* $(p_n)_{n=1}^{\infty}$*, by* p_{∞}*. If the given chain* $(p_n)_{n=1}^{\infty}$ *is a finite sequence, then* $p_{\infty} = p_N$*, for some* $N \in \mathbb{N}$*.*

Lemma 2.17. *Let* A *be a unital algebra and let* $p \in A_{pro}$*. Then* $px = xp$*, for all* $x \in pAp$*.*

Proof. Let $x \in pAp$. Then there exists an element $a \in A$ such that $x = pap$. Then

$$px = p(pap) = p^2 ap = pap = x$$

and

$$xp = (pap)p = pap^2 = pap = x.$$

∎

Lemma 2.18. *Let* A *be a unital algebra and let* $p, q \in A_{pro}$ *satisfying that* pAp *is a subalgebra of* qAq*. Then* $pq = qp = p$*.*

Proof. By assumption, $pAp \subset qAq$. Let $x \in pAp$. Then, by the previous lemma,

$$xq = qx.$$

Since $p \in pAp$, we can get that

$$pq = qp.$$

Now, observe that

$$pq = p^3 q = p\left(p^2 q\right) = p\left(p(pq)\right) = p\left(pqp\right) = p$$

since $pAp \subset qAq$. ∎

Corollary 2.19. *Let B be a unital algebra and A, an algebra over B. Let $(p_n)_{n=1}^{\infty} \subset A_{pro}$ be a chain of projections. Then, for any fixed $j < N \in \mathbb{N}$, $(p_j p_{j+1} ... p_N)\, A\, (p_j p_{j+1} ... p_N) = p_j A p_j$.*

Proof. By the previous lemmas, we have that

$$\begin{aligned}(p_j p_{j+1} ... p_N)\, A\, (p_j p_{j+1} ... p_N) &= p_j p_{j+1} ... p_N A p_N ... p_{j+1} p_j \\ &= p_j p_{j+1} ... (p_N A p_N) ... p_{j+1} p_j \\ &= p_j ... (p_{N-1} A p_{N-1}) ... p_j \\ &= p_j A p_j.\end{aligned}$$

∎

Definition 2.7. *Let B be a unital algebra and A, an algebra over B. Let (A, φ) be a NCPSpace over B, with its B-functional $\varphi : A \to B$. Suppose that we have a chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$ satisfying*

$$\mathbb{C} \subset^{\varphi_0} B \subset p_1 A p_1 \subset p_2 A p_2 \subset .... \subset p_\infty A p_\infty \subset A.$$

and

$$\varphi(p_k) \overset{denote}{=} \alpha_k \cdot 1_B \in \mathbb{C} \cdot 1_B, \text{ for all } k \in \mathbb{N}.$$

(It is possible that this sequence of projections is a finite sequence, i.e, $p_\infty = p_N$, for some N) We will call such chains of projections, scalar-valued chains of projections.

Suppose that we have a scalar-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$. We can naturally define conditional expectations

$$\varphi_{j+1} : p_{j+1} A p_{j+1} \to p_j A p_j$$

by

$$\varphi_{j+1}(p_{j+1} a p_{j+1}) \overset{def}{=} p_j a p_j,$$

for all $a \in A$, $j \in \mathbb{N}$. Notice that, by the previous lemmas,

$$\begin{aligned}\varphi_{j+1}(p_{j+1} a p_{j+1}) &= p_j (p_{j+1} a p_{j+1}) p_j \\ &= p_j p_{j+1} a p_{j+1} p_j \\ &= (p_j p_{j+1})\, a\, (p_j p_{j+1}) \\ &= p_j a p_j,\end{aligned}$$

for all $a \in A$. i.e,

$$\varphi_{j+1}(x) = p_j x p_j, \text{ for all } x \in p_{j+1} A p_{j+1},$$

for all $j \in \mathbb{N}$.

Hence we can have a tower of amalgamated NCPSpaces induced by a scalar-valued chain of projections, $(p_k)_{k=1}^{\infty} \subset A_{pro}$,

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} p_1Ap_1 \subset^{\varphi_2} p_2Ap_2 \subset^{\varphi_3} \cdots,$$

where a linear functional $\varphi_0 : B \to \mathbb{C}$ is arbitrary given.

Let B be a unital algebra and (A, φ), a NCPSpace over B. Suppose that we have a tower of amalgamated NCPSpaces induced by a scalar-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$ and the given linear functional $\varphi_0 : B \to \mathbb{C}$. If we define a map

$$E_{j+1} : p_{j+1}Ap_{j+1} \to \mathbb{C}$$

by

$$E_{j+1}(x) = \frac{1}{\alpha_{j+1}} \cdot \varphi_0\varphi_1...\varphi_{j+1}(x), \text{ for all } x \in A_{j+1},$$

then it is a well-defined linear functional. Recall that α_{j+1} is the scalar-part of $\varphi(p_{j+1})$.

Now, we will introduce compressed R-transform theory in [10] ; Let B be a unital algebra and (A, φ), a NCPSpace over B, with its B-functional $\varphi : A \to B$. Let $p \in A_{pro}$ such that

$$B \subset pAp \subset A$$

and

$$\varphi(p) \overset{denote}{=} b_0 \in C_A(B) \cap B_{inv}.$$

Note that if we have a scalar-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$, then every p_j satisfies

$$\varphi(p_j) = \alpha_j \cdot 1_B \in C_A(B) \cap B_{inv},$$

for all $j \in \mathbb{N}$ and $\alpha_j \in \mathbb{C}$.

We can define the conditional expectation

$$\varphi_p \overset{def}{=} b_0^{-1} \cdot \varphi \mid_{pAp}: pAp \to B.$$

So, we can define a NCPSpace over B, $\left(pAp, \, \varphi_p\right)$ and we call it a compressed NCPSpace by $p \in A_{pro}$. In [10], we observed the amalgamated R-transform theory on this compressed NCPSpace over B.

Definition 2.8. *Let B be a unital algebra and $b \in B$. Let (A, φ) be a NCPSpace over B. Define a symmetric cumulants of B-valued random variables $x_1, ..., x_s \in (A, \varphi)$ by*

$$k_n^{symm(b)}\left(x_{i_1}, ..., x_{i_n}\right) = c^{(n)}\left(x_{i_1} \otimes bx_{i_2} \otimes ... \otimes bx_{i_n}\right),$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. *Then we can define the symmetric R-transform of* $x_1, ..., x_s$, *by* $b \in B$, *by*

$$R_{x_1,...,x_s}^{symm(b)}(z_1, ..., z_s) = \textstyle\sum_{n=1}^{\infty} \displaystyle\sum_{i_1,...,i_n \in \{1,...,s\}} k_n^{symm(b)}(x_{i_1}, ..., x_{i_n})\, z_{i_1} ... z_{i_n},$$

as a B*-formal series in* Θ_B^s.

Theorem 2.20. *(See Theorem 2.1 in [10]) Let* (A, φ), $p \in A_{pro}$ *and* $b_0 \in B$ *be given as before. Let* $x_1, ..., x_s \in (A, \varphi)$ *be* B*-valued random variables* $(s \in \mathbb{N})$ *and assume that* $\{p\}$ *and* $\{x_1, ..., x_s\}$ *are free over* B. *Then*

$$R_{px_1p,...,px_sp}^{(\varphi_p)\,:\,t}(z_1, ..., z_s) = R_{x_1,...,x_s}^{symm(b_0)}(z_1, ..., z_s)$$

Proof. Fix $n \in \mathbb{N}$ and $(i_1, ..., i_n) \in \{1, ..., s\}^n$. Then

$$\begin{aligned} coef_{i_1,...,i_n}\left(R_{px_1p,...,px_sp}^{(\varphi_p)\,:\,t}\right) &= c_{\varphi_p}^{(n)}\left(px_{i_1}p \otimes px_{i_2}p \otimes ... \otimes px_{i_n}p\right) \\ &= \sum_{\pi \in NC(n)} \widehat{\varphi_p}(\pi)\left(px_{i_1}p \otimes ... \otimes px_{i_n}p\right)\mu(\pi, 1_n) \\ &= \sum_{\pi \in NC(n)} \widehat{b_0^{-1}\varphi}\,(\pi)\left(px_{i_1}p \otimes ... \otimes px_{i_n}p\right)\mu(\pi, 1_n) \end{aligned}$$

(2.20.1)

$$= \sum_{\pi \in NC'(n+1)} b_0^{-1}\, \widehat{\varphi_p}\,(\pi)\left(p \otimes x_{i_1}p \otimes x_{i_2}p \otimes ... \otimes x_{i_n}p\right)\mu(\pi, 1_{n+1}),$$

where $NC'(n+1) = \{\theta \in NC(n+1) : (1) \in \theta\}$ which is lattice-isomorphic to $NC(n)$. By the B-freeness of $\{p\}$ and $\{x_1, ..., x_s\}$, we have that (2.20.1) is equivalent to (2.20.2) (For this squivalence, see the proof of Theorem 2.1 in [10]) ;

(2.20.2)

$$\begin{aligned} &\sum_{\pi \in NC'(n+1)} b_0^{-1} \cdot \widehat{c_\varphi}\left(\pi \cup Kr(\pi)\right)\left(1_B \otimes p \otimes x_{i_1} \otimes p \otimes ... \otimes x_{i_n} \otimes p\right) \\ &\quad = \sum_{\pi \in NC'(n+1)} b_0^{-1} \cdot \widehat{c_\varphi}\,(\pi)\left(1_B \otimes x_{i_1} \otimes ... \otimes x_{i_n}\right) \cdot \widehat{c_{\varphi_p}}\left(Kr(\pi)\right)\left(p \otimes ... \otimes p\right) \end{aligned}$$

since p is B-central, in the sense of [9]

$$= \sum_{\pi \in NC'(n+1)} b_0^{-1} \cdot \widehat{c_\varphi}(\pi)\left(1_B \otimes x_{i_1} \otimes ... \otimes x_{i_n}\right) b_0^{|Kr(\pi)|}$$

$$= \sum_{\pi \in NC'(n+1)} b_0^{-1} \cdot \widehat{c_\varphi}(\pi) \left(1_B \otimes x_{i_1} \otimes ... \otimes x_{i_n}\right) b_0^{(n+1)-|\pi|},$$

since $|\pi|+|Kr(\pi)| = n+1$, for all $\pi \in NC(n)$. (Notice that $NC'(n+1) = NC(n)$.) Therefore,

$$k_n^{(\varphi_p)\,:\,t}\left(px_{i_1}p, ..., px_{i_n}p\right) = k_n^{(\varphi)\,:\,symm(b_0)}\left(x_{i_1}, ..., x_{i_n}\right).$$

∎

Remark that $p \in A_{pro}$ commutes with B. Since $b = pbp$, we have that

$$pb = p(pbp) = p^2bp = pbp = pbp^2 = (pbp)p = bp,$$

for all $b \in B$. i.e, $p \in (A, \varphi)$ is B-central, in the sense of [9].

Let $(p_k)_{k=1}^{\infty} \subset A_{pro}$ be a scalar-valued chain of projections in (A, φ). Then each $p_j \in A_{pro}$ satisfies $\varphi(p_j) = \alpha_j \cdot 1_B$, for some $\alpha_j \in \mathbb{C}$. Therefore, each $\varphi(p_j) \in C_{A_j}(A_{j-1}) \cap (A_{j-1})_{inv}$, for all $j \in \mathbb{N}$, with $A_0 = B$.

Corollary 2.21. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Let $x_1, ..., x_s \in (A, \varphi)$ be B-valued random variables ($s \in \mathbb{N}$). Suppose that we have $p \in A_{pro}$ such that $B \subset pAp \subset A$ and $\varphi(p) \overset{denote}{=} b_0 \in C_A(B) \cap B_{inv}$. If $\{p\}$ and $\{x_1, ..., x_s\}$ are free over B, then*

$$\begin{aligned} k_n^{(\varphi_p)}\left(px_{i_1}p, ..., px_{i_n}p\right) &:= c_{\varphi_p}^{(n)}\left(px_{i_1}p \otimes b_{i_2}(px_{i_2}p) \otimes ... \otimes b_{i_n}(px_{i_n}p)\right) \\ &= c_{\varphi}^{(n)}\left(x_{i_1} \otimes b_0 b_{i_2} x_{i_2} \otimes ... \otimes b_0 b_{i_n} x_{i_n}\right), \end{aligned}$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$, where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary.

Proof. The proof is similar to the proof of the previous theorem. Fix $n \in \mathbb{N}$ and $(i_1, ..., i_n) \in \{1, ..., s\}^n$. Observe that

$$k_n^{(\varphi_p)}\left(px_{i_1}p, ..., px_{i_n}p\right) = c_{\varphi_p}^{(n)}\left(px_{i_1}p \otimes b_{i_2}px_{i_2}p \otimes ... \otimes pb_{i_n}x_{i_n}p\right)$$

where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary. Since $pb_{i_j}p = b_{i_j}$, we have that

$$pb_{i_j} = p\left(pb_{i_j}p\right) = p^2b_{i_j}p = pb_{i_j}p = pb_{i_j}p^2 = (pb_{i_j}p)p = b_{i_j}p,$$

for all $j = 1, ..., n$. So, we have that

$$k_n^{(\varphi_p)}(px_{i_1}p, ..., px_{i_n}p) = c_{\varphi_p}^{(n)}\left(px_{i_1}p \otimes px'_{i_2}p \otimes ... \otimes px'_{i_n}p\right),$$

where $x'_{i_j} = b_{i_j}x_{i_j}$, for all $j = 1, ..., n$. Similar to the proof of the previous theorem, we have that

$$c_{\varphi_p}^{(n)}\left(px_{i_1}p \otimes px'_{i_2}p \otimes ... \otimes px'_{i_n}p\right) = c_{\varphi}^{(n)}\left(x_{i_1} \otimes b_0x'_{i_2} \otimes ... \otimes b_0x'_{i_n}\right),$$

where $b_0 = \varphi(p) \in C_A(B) \cap B_{inv}$. Therefore,

$$k_n^{(\varphi_p)}(x_{i_1}, ..., x_{i_n}) = c_{\varphi}^{(n)}\left(x_{i_1} \otimes b_0x'_{i_2} \otimes ... \otimes b_0x'_{i_n}\right).$$

∎

By the previous corollary, we can get the following theorem ;

Theorem 2.22. *(Also see Theorem 2.5 and Corollary 2.7 in [10]) Let B be a unital algebra and let (A, φ) be a NCPSpace over B. Let $p \in A_{pro}$ satisfy $\varphi(p) \in C_A(B) \cap B_{inv}$ and suppose that $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ are two subsets of B-valued random variables in (A, φ). Assume that $\{p\}$ and $X \cup Y$ are free over B, in (A, φ). If X and Y are free over B, in (A, φ), then pXp and pYp are free over B, in $\left(pAp, \varphi_p\right)$.*

Proof. Suppose that $\{p\}$ and $X \cup Y$ are free over B. Then, by the previous corollary, we can get that

$$k_n^{(\varphi_p)}(a_{i_1}, ..., a_{i_n}) = c_{\varphi_p}^{(n)}(pa_{i_1}p \otimes b_{i_2}pa_{i_2}p \otimes ... \otimes b_{i_n}pa_{i_n}p)$$

$$= c_{\varphi_p}^{(n)}(pa_{i_1}p \otimes pb_{i_2}a_{i_2}p \otimes ... \otimes pb_{i_n}a_{i_n}p)$$

(2.22.1)
$$= c_{\varphi}^{(n)}(a_{i_1} \otimes b_0(b_{i_2}a_{i_2}) \otimes ... \otimes b_0(b_{i_n}a_{i_n}))$$

where $b_0 = \varphi(p) \in C_A(B) \cap B_{inv}$, $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary and where $a_{i_1}, ..., a_{i_n} \in X \cup Y$. We can rewrite (2.22.1) as

$$c_{\varphi}^{(n)}\left(a_{i_1} \otimes b'_{i_2}a_{i_2} \otimes ... \otimes b'_{i_n}a_{i_n}\right), \tag{2.22.2}$$

where $b'_{i_2}, ..., b'_{i_n} \in B$ are arbitrary such that $b'_{i_j} = b_0b_{i_j}$, for all $j = 1, ..., n$. Since X and Y are free over B, in (A, φ), we have that

$$(2.22.2) \quad = \begin{cases} c_{\varphi}^{(n)}\left(x_{i_1} \otimes b'_{i_2} x_{i_2} \otimes ... \otimes b'_{i_n} x_{i_n}\right) & \text{or} \\ \\ c_{\varphi}^{(n)}\left(y_{i_1} \otimes b'_{i_2} y_{i_2} \otimes ... \otimes b'_{i_n} y_{i_n}\right). \end{cases}$$

$$= \begin{cases} k_n^{(\varphi_p)}\left(px_{i_1}p, ..., px_{i_n}p\right) & \text{or} \\ \\ k_n^{(\varphi_p)}\left(py_{i_1}p, ..., py_{i_n}p\right). \end{cases}$$

Therefore, pXp and pYp are also free over B, in $\left(pAp,\, \varphi_p\right)$. ∎

Corollary 2.23. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Let $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ be subsets of B-valued random variables in (A, φ). Assume that X and Y are free over B, in (A, φ). Now, let $p \in A_{pro}$ such that $\{p\}$ and $X \cup Y$ are free over B, in (A, φ). By $b_0 \in C_A(B) \cap B_{inv}$, we will denote $\varphi(p) \in C_A(B) \cap B_{inv}$. Then*

(1) $R_{px_1p,...,px_sp}^{(\varphi_p)\,:\,t}(z_1, ..., z_s) = R_{x_1,...,x_s}^{(\varphi)\,:\,symm(b_0)}(z_1, ..., z_s)$ *and*

$$R_{px_1p,...,px_sp}^{(\varphi_p)}(z_1, ..., z_s) = \sum_{n=1}^{\infty} \sum_{i_1,...,i_n \in \{1,...,s\}} c_{\varphi}^{(n)}\left(x_{i_1} \otimes b_0 b_{i_2} x_{i_2} \otimes ... \otimes b_0 b_{i_n} x_{i_n}\right) z_{i_1} ... z_{i_n},$$

where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary.

(2) $R_{px_1p,...,px_sp,py_1p,...,py_sp}^{(\varphi_p)}(z_1, ..., z_{2s})$

$$= R_{px_1p,...,px_sp}^{(\varphi_p)}(z_1, ..., z_s) + R_{py_1p,...,py_sp}^{(\varphi_p)}(z_{s+1}, ..., z_{2s}).$$

(3) $R_{px_1p+py_1p,...,px_sp+py_1p}^{(\varphi_p)}(z_1, ..., z_s)$

$$= \left(R_{px_1p,...,px_sp}^{(\varphi_p)} + R_{py_1p,...,py_sp}^{(\varphi_p)}\right)(z_1, ..., z_s).$$

(4) $R_{(px_1p)(py_1p),...(px_sp)(py_sp)}^{(\varphi_p)}(z_1, ..., z_s)$

$$= \left(R_{px_1p,...,px_sp}^{(\varphi_p)} \boxed{*}_B\, R_{py_1p,...,py_sp}^{(\varphi_p)\,:\,t}\right)(z_1, ..., z_s)$$

$$= \left(R_{px_1p,...,px_sp}^{(\varphi_p)} \boxed{*}_B\, R_{y_1,...,y_s}^{(\varphi)\,:\,t}\right)(z_1, ..., z_s)$$

Proof. (1), (2) and (3) are proved in the previous theorems and corollary. Since pXp and pYp are free over B, in $\left(pAp, \varphi_p\right)$, we have that

$$R_{(px_1p)(py_1p),...,(px_sp)(py_sp)}^{(\varphi_p)} = R_{px_1p,...,px_sp}^{(\varphi_p)} \boxed{*}_B\, R_{py_1p,...,py_sp}^{(\varphi_p)}.$$

Also, we can get the following result ; now, fix $n \in \mathbb{N}$ and $(i_1, ..., i_n) \in \{1, ..., s\}^n$.

$$
\begin{aligned}
& coef_{i_1,...,i_n}\left(R^{(\varphi_p)}_{(px_1p)(py_1p),...,(px_sp)(py_1p)}\right) \\
& \quad = c^{(n)}_{\varphi_p}\left(px_{i_1}y_{i_1}p \otimes b_{i_2}px_{i_2}y_{i_2}p \otimes ... \otimes b_{i_n}px_{i_n}y_{i_n}p\right),
\end{aligned}
$$

since $(px_{i_j}p)(py_{i_j}p) = px_{i_j}p^2y_{i_j}p = px_{i_j}py_{i_j}p = px_{i_j}y_{i_j}p$, for all $j = 1, ..., n$, where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary

$$
\begin{aligned}
& = c^{(n)}_{\varphi}\left(x_{i_1}y_{i_1} \otimes b_0b_{i_2}x_{i_2}y_{i_2} \otimes ... \otimes b_0b_{i_n}x_{i_n}y_{i_n}\right) \\
& = c^{(n)}_{\varphi}\left(x_{i_1}y_{i_1} \otimes b'_{i_2}x_{i_2}y_{i_2} \otimes ... \otimes b'_{i_n}x_{i_n}y_{i_n}\right)
\end{aligned}
$$

where $b'_{i_j} = b_0b_{i_j} \in B$, for all $j = 1, ..., n$

$$
= \sum_{\pi \in NC(n)} \widehat{c_{\varphi}}\left(\pi \cup Kr(\pi)\right)\left(x_{i_1} \otimes y_{i_1} \otimes b'_{i_2}x_{i_2} \otimes y_{i_2} \otimes ... \otimes b'_{i_n}x_{i_n} \otimes y_{i_n}\right)
$$

by the freeness of $\{x_1, ..., x_s\}$ and $\{y_1, ..., y_s\}$. So,

$$
coef_{i_1,..,i_n}\left(R^{(\varphi_p)}_{(px_1p)(py_1p),...,(px_sp)(py_sp)}\right) = coef_{i_1,...,i_n}\left(R^{(\varphi_p)}_{px_1p,...,px_sp} \boxed{*}_B \, R^{(\varphi)\,:\,t}_{y_1,...,y_s}\right).
$$

$\blacksquare$

Notations Let B be a unital algebra and (A, φ), a NCPSpace over B. Now consider the following tower of amalgamated NCPSpaces,

$$
\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} p_1Ap_1 \subset^{\varphi_2} p_2Ap_2 \subset^{\varphi_3} \cdots,
$$

induced by a scalar-valued chain of projections in A, $(p_k)_{k=1}^{\infty} \subset A_{pro}$, satisfying $\varphi(p_k) = \alpha_k \cdot 1_B \in C_A(B) \cap B_{inv}$, $\forall k$, and the given linear functional $\varphi_0 : B \to \mathbb{C}$. As we defined before,

$$
\varphi_{j+1}(p_{j+1}ap_{j+1}) = p_jap_j, \text{ for each } j \in \mathbb{N}, \text{ for all } a \in A.
$$

As in Section 2.1, we can apply the R-transform theory on a tower of amalgamated NCPSpaces to this compressed amalgamated NCPSpaces induced by a scalar-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$ and the given linear functional φ_0.

Theorem 2.24. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Let $(p_k)_{k=1}^{\infty} \subset A_{pro}$ be a scalar-valued chain of projections. Consider the tower of amalgamated NCPSpaces induced by the chain of projections $(p_k)_{k=1}^{\infty}$ and a linear functional $\varphi_0 : B \to \mathbb{C}$. Then, for any $j \in \mathbb{N}$, if two subsets of B-valued random* *variables X and Y are free over B, in (A, φ), and if p_∞ and $X \cup Y$ are free over B, in (A, φ), then $p_j X p_j$ and $p_j Y p_j$ are free over B, in $\left(p_j A p_j,\, \frac{1}{\alpha_1} \cdot E_{1,\,j}\right)$.* $\square$

Notice that

$$\frac{1}{\alpha_1} = \frac{1}{E_{1,\,j}(p_1 \ldots p_j)} = \frac{1}{E_{1,\,j}(p_1)} = \frac{1}{\varphi_1 \ldots \varphi_j(p_1)} = \frac{1}{\varphi_1(p_1)} = \frac{1}{\varphi(p_1)}.$$

As an application of of the above theorem, we can get the following R-transform calculus ;

Theorem 2.25. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Let $(p_k)_{k=1}^{\infty} \subset A_{pro}$ be a scalar-valued chain of projections. Suppose that we have a tower of amalgamated NCPSpaces induced by the chain of projections and a linear functional $\varphi_0 : B \to \mathbb{C}$. Let $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ be two B-free subsets in (A, φ). If p_∞ and $X \cup Y$ are free over B, in (A, φ), then, for any fixed $j \in \mathbb{N}$, we have that*

(1) $R_{p_j x_1 p_j, \ldots, p_j x_s p_j, p_j y_1 p_j, \ldots, p_j y_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_1, ..., z_{2s})$

$$= R_{p_j x_1 p_j, \ldots, p_j x_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_1, ..., z_s) + R_{p_j y p_j, \ldots, p_j y_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_{s+1}, ..., z_{2s}).$$

(2) $R_{p_j x_1 p_j + p_j y_1 p_j, \ldots, p_j x_s p_j + p_j y_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_1, ..., z_{2s})$

$$= R_{p_j x_1 p_j, \ldots, p_j x_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_1, ..., z_s) + R_{p_j y p_j, \ldots, p_j y_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_1, ..., z_s).$$

(3) $R_{(p_j x_1 p_j)(p_j y_1 p_j), \ldots, (p_j x_s p_j)(p_j y_s p_j)}^{(\frac{1}{\alpha_1} E_{1,\,j})}(z_1, ..., z_s)$

$$= \left(R_{p_j x_1 p_j, \ldots, p_j x_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})} \ \boxed{*}_B \ R_{p_j y_1 p_j, \ldots, p_j y_s p_j}^{(\frac{1}{\alpha_1} E_{1,j})\,:\,t} \right)(z_1, ..., z_s)$$

$$= \left(R_{p_j x_1 p_j, \ldots, p_j x_s p_j}^{(\frac{1}{\alpha_1} E_{1,\,j})} \ \boxed{*}_B \ R_{y_1, \ . \ . \ . \ , y_s}^{(\varphi)\,:\,t} \right)(z_1, ..., z_s).$$

$\square$

By using the same idea of the previous two theorems, we have that ;

Theorem 2.26. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Let $(p_k)_{k=1}^{\infty} \subset A_{pro}$ be a scalar-valued chain of projections. Consider the tower of amalgamated NCPSpaces induced by the chain of projections $(p_k)_{k=1}^{\infty}$ and a linear functional $\varphi_0 : B \to \mathbb{C}$. Then, for any $k < j \in \mathbb{N}$, if two subsets of B-valued random variables X and Y are free over A_k, in (A, φ), then $p_j X p_j$ and $p_j Y p_j$ are free over B, in $\left(p_j A p_j,\, \frac{1}{\alpha_{k+1}} \cdot E_{k+1,\,j}\right)$.* $\square$

Theorem 2.27. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Let $(p_k)_{k=1}^{\infty} \subset A_{pro}$ be a scalar-valued chain of projections. Suppose that we have a tower of amalgamated NCPSpaces induced by the chain of projections and a linear functional $\varphi_0 : B \to \mathbb{C}$. Let $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ be two A_k-free subsets in (A, φ). Then, for any fixed $k < j \in \mathbb{N}$, we have that*

(1) $$R_{p_j x_1 p_j, ..., p_j x_s p_j, p_j y_1 p_j, ..., p_j y_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_1, ..., z_{2s})$$
$$= R_{p_j x_1 p_j, ..., p_j x_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_1, ..., z_s) + R_{p_j y p_j, ..., p_j y_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_{s+1}, ..., z_{2s}).$$

(2) $$R_{p_j x_1 p_j + p_j y_1 p_j, ..., p_j x_s p_j + p_j y_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_1, ..., z_{2s})$$
$$= R_{p_j x_1 p_j, ..., p_j x_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_1, ..., z_s) + R_{p_j y p_j, ..., p_j y_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_1, ..., z_s).$$

(3) $$R_{(p_j x_1 p_j)(p_j y_1 p_j), ..., (p_j x_s p_j)(p_j y_s p_j)}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})}(z_1, ..., z_s)$$
$$= \left(R_{p_j x_1 p_j, ..., p_j x_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})} \ \boxed{*}_B \ R_{p_j y_1 p_j, ..., p_j y_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1, j})\,:\,t} \right)(z_1, ..., z_s)$$
$$= \left(R_{p_j x_1 p_j, ..., p_j x_s p_j}^{(\frac{1}{\alpha_{k+1}} E_{k+1,\, j})} \ \boxed{*}_B \ R_{y_1, \ . \ . \ . \ , y_s}^{(\varphi)\,:\,t} \right)(z_1, ..., z_s).$$

□

3. Compatibility of towers of Amalgamated Noncommutative Probability Spaces

Throughout this chapter, we will consider the tower of amalgamated NCPSpaces,

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset \cdots.$$

In this chapter, we will consider the compatibility of amalgamated NCPSpaces.

Definition 3.1. *Let (A, φ) be a NCPSpace over B and let (A, φ_0) be a NCPSpace (over $\mathbb{C}$). We say that (A, φ) and (A, φ_0) are compatible if*

$$\varphi_0(x) = \varphi_0\left(\varphi(x)\right), \ \textit{for all } x \in A.$$

Now let D be a unital subalgebra of A such that $1_D = 1_A (= 1_B)$ and assume that there is a conditional expectation $\varphi' : A \to D$ and hence we have a NCPSpace *over* D, (A, φ'). *(i.e,* $B \subset D \subset A$.*) We say that two amalgamated NCPSpaces* $(A, \varphi' : A \to D)$ *and* $(A, \varphi : A \to B)$ *are compatible if*

$$\varphi(x) = \varphi\left(\varphi'(x)\right), \text{ for all } x \in A.$$

Proposition 3.1. *Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} \cdots$$

be a tower of amalgamated NCPSpaces. Then, for any fixed $j \in \mathbb{N}$, $\left(A_{j+1}, \varphi_{j+1}\right)$ *and* $\left(A_{j+1}, \varphi_j \varphi_{j+1}\right)$ *are compatible.*

Proof. For any $x \in A_{j+1}$, we have that

$$\varphi_j \varphi_{j+1}(x) = \varphi_j \varphi_{j+1}^2(x) = \varphi_j \varphi_{j+1}\left(\varphi_{j+1}(x)\right),$$

since $\varphi_{j+1}^2(x) = \varphi_{j+1}\left(\varphi_{j+1}(x)\right) = \varphi_{j+1}(x)$. ∎

In general, we can get the following facts ;

Theorem 3.2. *Let* B *be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} \cdots$$

be a tower of amalgamated NCPSpaces. For any fixed $k < i < j$ *in* $\mathbb{N}$, $(A_j,\ E_{i,j})$ *and* $(A_j,\ E_{k,j})$ *are compatible.*

Proof. For any $x \in A_j$, we have that

$$\begin{aligned} E_{k,j}(x) &= \varphi_k ... \varphi_i ... \varphi_j(x) = \varphi_k ... \varphi_{i-1}\left(\varphi_i ... \varphi_j(x)\right) \\ &= E_{k,\, i-1}\left(E_{i,j}(x)\right) = E_{k,\, i-1}\left(E_{i,j}^2(x)\right) \end{aligned}$$

since $E_{i,j} : A_j \to A_{i-1}$ is a conditional expectation

$$= E_{k,\, i-1} E_{i,j}\left(E_{i,j}(x)\right) = E_{k,j}\left(E_{i,j}(x)\right).$$

Therefore,

$$E_{k,j}(x) = E_{k,j}\left(E_{i,j}(x)\right) \in A_{k-1}, \text{ for all } x \in A_j.$$

∎

Theorem 3.3. *Let B be a unital algebra and suppose that we have a tower of amalgamated NCPSpaces as before. Then, for any fixed $k < j$ in $\mathbb{N}$, $(A_j,\, E_j)$ and $(A_j,\, E_{k,\, j})$ are compatible.*

Proof. For any $x \in A_j$, we have that

$$\begin{aligned} E_j(x) &= \varphi_0\varphi_1..\varphi_k...\varphi_j(x) = E_{k-1}\left(E_{k,\, j}(x)\right) \\ &= E_{k-1}\left(E_{k,\, j}^2(x)\right) = E_{k-1}E_{k,\, j}\left(E_{k,\, j}(x)\right) \\ &= E_j\left(E_{k,\, j}(x)\right) \in \mathbb{C}. \end{aligned}$$

▮

The above two theorems shows us that we can freely use the compatibility on the tower.

Lemma 3.4. *(See [18] and [12]) Let B be a unital algebra and A, an algebra over B. Suppose a NCPSpace (A, φ_0) and a NCPSpace over B, (A, φ) are compatible. Let $x_1, ..., x_s \in A$ be operators such that $\{x_1, ..., x_s\}$ and B are free in (A, φ_0). Then*

$$\begin{aligned} k_n^{(\varphi)}\left(x_{i_1}, ..., x_{i_n}\right) &= c^{(n)}\left(x_{i_1} \otimes b_{i_2}x_{i_2} \otimes ... \otimes b_{i_n}x_{i_n}\right) \\ &= \left(\varphi_0(b_{i_2})...\varphi_0(b_{i_n})\right) \cdot k_n^{(\varphi_0)}\left(x_{i_1}, ..., x_{i_n}\right) \cdot 1_B, \end{aligned}$$

in B, where $k_n^{(\varphi_0)}(...)$ is the scalar-valued cumulants of $x_1, ..., x_s$, in the sense of Speicher and Nica, and where $b_{i_2}, ..., b_{i_n} \in B$ are arbitrary, for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. □

In the above lemma, $k_n^{(\varphi_0)}(...)$ is our $k_n^{(\varphi_0)\,:\,t}(...)$. (See Chapter 1 and [6], [7])

Proposition 3.5. *Let B be a unital algebra and let*

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} \cdots$$

be a tower of amalgamated NCPSpaces. Fix $k < j \in \mathbb{N}$. Suppose that $X = \{x_1, ..., x_s\}$, $Y = \{y_1, ..., y_s\} \subset A_{j+1}$ are two subsets of operators and assume that $X \cup Y$ is free from A_k, in $(A_{j+1},\, E_{j+1})$. If X and Y are free in $(A_{j+1},\, E_{j+1})$, then X and Y are free over A_k, in $(A_{j+1},\, E_{k+1,\, j+1})$.

Proof. Now take the following (sub)tower of the given tower

$$\mathbb{C} \subset^{E_k} A_k \subset^{E_{k+1,\, j+1}} A_{j+1}.$$

Then, by the previous theorems, $(A_{j+1},\, E_{j+1})$ and $(A_{j+1},\, E_{k+1,\,j+1})$ are compatible. By hypothesis, for any $p_1, ..., p_n \in X \cup Y$, we have that

$$k_n^{(E_{k+1,\,j+1})}(p_1, ..., p_n) = c_{E_{k+1,\,j+1}}^{(n)}(p_1 \otimes b_2 p_2 \otimes ... \otimes b_n p_n)$$

where $b_2, ..., b_n \in B$ are arbitrary

$$= (E_{j+1}(b_2) \cdots E_{j+1}(b_n)) \cdot k_n^{(E_{j+1})}(p_1, ..., p_n) \cdot 1_B, \tag{3.5.1}$$

by the previous lemma, where $k_n^{(E_{j+1})}(...)$ is the scalar-valued cumulant, in the sense of Speicher and Nica. By the freeness of X and Y in $(A_{j+1},\, E_{j+1})$, (3.5.1) goes to

$$= \begin{cases} (E_{j+1}(b_2) \cdots E_{j+1}(b_n)) \cdot k_n^{(E_{j+1})}(x_{i_1}, ..., x_{i_n}) \cdot 1_B & \text{or} \\ (E_{j+1}(b_2) \cdots E_{j+1}(b_n)) \cdot k_n^{(E_{j+1})}(y_{i_1}, ..., y_{i_n}) \cdot 1_B & \end{cases}$$

$$= \begin{cases} k_n^{(\varphi)}(x_{i_1}, ..., x_{i_n}) & \text{or} \\ k_n^{(\varphi)}(y_{i_1}, ..., y_{i_n}), & \end{cases}$$

for all $(i_1, ..., i_n) \in \{1, ..., s\}^n$, $n \in \mathbb{N}$. Therefore, X and Y are free over A_k, in $(A_{j+1},\, E_{k+1,\,j+1})$. ▮

Proposition 3.6. *Let B be a unital algebra and suppose that we have a tower of amalgamated NCPSpaces as before. Fix $k < j$ in $\mathbb{N}$. Let $X = \{x_1, ..., x_s\}$ and $Y = \{y_1, ..., y_s\}$ be two subsets of operators in A_{j+1}. Assume that $X \cup Y$ is free from A_k in $(A_{j+1},\, E_{j+1})$. If X and Y are free in $(A_{j+1},\, E_{j+1})$, then*

(1) $R_{x_1,...,x_s,y_1,...,y_s}^{(E_{k+1,\,j})}(z_1, ..., z_s) = R_{x_1,...,x_s}^{(E_{k+1,\,j})}(z_1, ..., z_s) + R_{y_1,...,y_s}^{(E_{k+1,\,j})}(z_{s+1}, ..., z_{2s})$.

(2) $R_{x_1+y_1,...,x_s+y_s}^{(E_{k+1,\,j})}(z_1, ..., z_s) = \left(R_{x_1,...,x_s}^{(E_{k+1,\,j})} + R_{y_1,...,y_s}^{(E_{k+1,\,j})}\right)(z_1, ..., z_s)$.

(3) $R_{x_1y_1,...,x_sy_s}^{(E_{k+1,\,j})}(z_1, ..., z_s) = \left(R_{x_1,...,x_s}^{(E_{k+1,\,j})} \;\boxed{*}_{A_k}\; R_{y_1,...,y_s}^{(E_{k+1,\,j})}\right)(z_1, ..., z_s)$.

(Recall that $E_{k+1,\,k+1} = \varphi_{k+1} : A_{k+1} \rightarrow A_k$, for all k.)

Proof. Remember that $(A_{j+1},\, E_{j+1})$ and $(A_{j+1},\, E_{k+1,\,j})$ are compatible, for any $k < j$ in $\mathbb{N}$. Since $X \cup Y$ and A_k are free in $(A_{j+1},\, E_{j+1})$, if X and Y are free in $(A_{j+1},\, E_{j+1})$, then X and Y are free over A_k, in $(A_{j+1},\, E_{k+1,\,j})$, by the previous proposition. ▮

4. Ladders of Amalgamated Noncommutative Probability Spaces

In this chapter, we will consider new algebraic structure, so-called a ladder of amalgamated NCPSpaces. Suppose that we have two towers of amalgamated NCPSpaces,

$$\mathbb{C} \subset^{\varphi_0} B \subset^{\varphi_1} A_1 \subset^{\varphi_2} A_2 \subset^{\varphi_3} A_3 \subset^{\varphi_4} A_4 \subset^{\varphi_5} \cdots$$

and

$$\mathbb{C} \subset^{\varphi_0'} B' \subset^{\varphi_1'} A_1' \subset^{\varphi_2'} A_2' \subset^{\varphi_3'} A_3' \subset^{\varphi_4'} A_4' \subset^{\varphi_5'} \cdots.$$

Assume that there is a relation between this two towers ;

$$A_j' \subset A_j \text{ is a subalgebra, for all } j \in \mathbb{N} \cup \{0\},$$

with $1_{A_j'} = 1_{A_j}$, where $A_0' = B'$ and $A_0 = B$. Then, for any k, we have the following square as a part of the ladder,

$$\begin{array}{ccc} A_k & \subset^{E_{k+1,\,j}} & A_j \\ \cup^{i_k} & & \cup^{i_j} \\ A_k' & \subset^{E_{k+1,\,j}'} & A_j', \end{array}$$

where $E_{k+1,\,j}' = \varphi_{k+1}...\varphi_j \mid_{A_j'}: A_j' \to A_k'$ is a conditional expectation and $i_k : A_k \to A_k'$, $i_j : A_j \to A_j'$ are conditional expectations. i.e, we have the following ladder generated by those two towers of amalgamated NCPSpaces,

$$\begin{array}{cccccccccccc} \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & A_1 & \subset^{\varphi_2} & A_2 & \subset^{\varphi_3} & A_3 & \subset^{\varphi_4} & \cdots \\ & & \cup^{i_0} & & \cup^{i_1} & & \cup^{i_2} & & \cup^{i_3} & & \cdots \\ \mathbb{C} & \subset^{\varphi_0'} & B' & \subset^{\varphi_1'} & A_1' & \subset^{\varphi_2'} & A_2' & \subset^{\varphi_3'} & A_3' & \subset^{\varphi_4'} & \cdots \end{array}$$

Notice that $i_k\varphi_{k+1} : A_{k+1} \to A_k'$ is a conditional expectation, for all $k \in \mathbb{N}$. Indeed,

$$i_k\varphi_{k+1}(a) = i_k(a) = a, \text{ for all } a \in A_k',$$

since $a \in A_k' \subset A_k$. Also,

$$i_k\varphi_{k+1}\left(axa'\right) = i_k\left(a\varphi_{k+1}(x)a'\right) = a\left(i_k\varphi_{k+1}(x)\right)a',$$

for all $a, a' \in A_k'$ and $x \in A_{k+1}$.

Definition 4.1. *We say that the following inclusions*

$$\begin{array}{ccccccccccc} \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & A_1 & \subset^{\varphi_2} & A_2 & \subset^{\varphi_3} & A_3 & \subset^{\varphi_4} & \cdots \\ & & \cup^{i_0} & & \cup^{i_1} & & \cup^{i_2} & & \cup^{i_3} & & \cdots \\ \mathbb{C} & \subset^{\varphi_0'} & B' & \subset^{\varphi_1'} & A_1' & \subset^{\varphi_2'} & A_2' & \subset^{\varphi_3'} & A_3' & \subset^{\varphi_4'} & \cdots \end{array}$$

is a commuting ladder of amalgamated NCPSpaces if conditions (i) and (ii) are satisfied ;

(i) Two towers of inclusions

$$\mathbb{C} \quad \subset^{\varphi_0} \quad B \quad \subset^{\varphi_1} \quad A_1 \quad \subset^{\varphi_2} \quad A_2 \quad \subset^{\varphi_3} \quad A_3 \quad \subset^{\varphi_4} \quad \cdots,$$

$$\mathbb{C} \quad \subset^{\varphi_0'} \quad B' \quad \subset^{\varphi_1'} \quad A_1' \quad \subset^{\varphi_2'} \quad A_2' \quad \subset^{\varphi_3'} \quad A_3' \quad \subset^{\varphi_4'} \quad \cdots$$

are towers of amalgamated NCPSpaces.

(ii) For any choice of a square of amalgamated NCPSaces,

$$\begin{array}{ccc} A_k & \subset^{\varphi_{k+1}} & A_{k+1} \\ \cup^{i_k} & & \cup^{i_{k+1}} \\ A_k' & \subset^{\varphi_{k+1}'} & A_{k+1}' \end{array}$$

is a commuting square of amalgamated NCPSpaces, i.e conditional expectations on that square satisfy that

$$i_k \varphi_{k+1} = \varphi_{k+1}' i_{k+1} : A_{k+1} \to A_k'.$$

Example 4.1. *Let B be a unital algebra and (A, φ), a NCPSpace over B. Suppose that we have a linear functional $\varphi_0 : B \to \mathbb{C}$. Also assume that there is a scalar-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$. Suppose that there is a NCPSpace over B, (D, ψ) such that $D \subset A$ and there is a coditional expectation $E_D^A : A \to D$ defined by $E_D^A(x) = x$, if $x \in D \subset A$ and $E_D^A(x) = 0_D$, otherwise. Then we can get a commuting ladder of (compressed) amalgamated NCPSpaces,*

$$\begin{array}{ccccccccc} \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & p_1 A p_1 & \subset^{\varphi_2} & p_2 A p_2 & \subset^{\varphi_3} & \cdots \\ & & \cup^{E_D^A} & & \cup^{E_D^A} & & \cup^{E_D^A} & & \\ \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & p_1 D p_1 & \subset^{\varphi_2} & p_2 D p_2 & \subset^{\varphi_3} & \cdots, \end{array}$$

where $E_D^A = E_D^A \mid_{p_j A p_j}$ *and* φ_j*'s are defined in Section 2.3 (i.e,* $\varphi_{j+1}(p_{j+1} a p_{j+1}) = p_j a p_j$*,* $\forall a \in A$*), for all* $j \in \mathbb{N}$*. Indeed, for any* j*, we have a commuting square of amalgamated NCPSpaces,*

$$\begin{array}{ccc} p_j A p_j & \subset^{\varphi_{j+1}} & p_{j+1} A p_{j+1} \\ \cup^{E_D^A} & & \cup^{E_D^A} \\ p_j D p_j & \subset^{\varphi_{j+1}} & p_{j+1} D p_{j+1}, \end{array}$$

since

$$\begin{aligned} E_D^A \ \varphi_{j+1}\left(p_{j+1} a p_{j+1}\right) &= E_D^A\left(p_j a p_j\right) \\ &= \begin{cases} p_j a p_j & \textit{if } a \in D \\ 0_D & \textit{otherwise.} \end{cases} \end{aligned}$$

and

$$\begin{aligned} \varphi_{j+1} \ E_D^A\left(p_{j+1} a p_{j+1}\right) &= \begin{cases} \varphi_{j+1}\left(p_{j+1} a p_{j+1}\right) & \textit{if } a \in D \\ \varphi_{j+1}\left(0_D\right) & \textit{otherwise} \end{cases} \\ &= \begin{cases} p_j a p_j & \textit{if } a \in D \\ 0_D & \textit{otherwise.} \end{cases} \end{aligned}$$

Hence the given ladder satisfies the condition (ii) of Definition 4.1. Since two towers in the ladder are towers of compressed amalgamated NCPSPaces, our ladder satisfies the condition (i) of Definition 4.1. Thus the ladder is a commuting ladder of amalgamated NCPSpaces.

Proposition 4.1. *Let*

$$\begin{array}{cccccccccccc} \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & A_1 & \subset^{\varphi_2} & A_2 & \subset^{\varphi_3} & A_3 & \subset^{\varphi_4} & \cdots \\ & & \cup^{i_0} & & \cup^{i_1} & & \cup^{i_2} & & \cup^{i_3} & & \cdots \\ \mathbb{C} & \subset^{\varphi_0'} & B' & \subset^{\varphi_1'} & A_1' & \subset^{\varphi_2'} & A_2' & \subset^{\varphi_3'} & A_3' & \subset^{\varphi_4'} & \cdots \end{array}$$

be a commuting ladder of amalgamated NCPSpaces. Then, for any $k < j$ *in* $\mathbb{N}$*, a square of inclusions*

$$\begin{array}{ccc} A_k & \subset^{E_{k+1,\,j}} & A_j \\ \cup^{i_k} & & \cup^{i_j} \\ A_k' & \subset^{E_{k+1,\,j}'} & A_j' \end{array}$$

is a commuting square of amalgamated NCPSpaces.

Proof. Fix $k < j$. By the definition of commuting squares of amalgamated NCPSpaces, it suffices to show that

$$i_k E_{k+1,\,j} = E'_{k+1,\,j} i_j : A_j \to A'_k$$

as a conditional expectation on A_j over A_k. Observe that, for any $x \in A_j$,

$$\begin{aligned}
i_k E_{k+1,\,j}(x) \quad & = i_k \varphi_{k+1} ... \varphi_j(x) = \left(i_k \varphi_{k+1}\right) \varphi_{k+2} ... \varphi_j(x) \\
& = \left(\varphi'_{k+1} i_{k+1}\right) \varphi_{k+2} ... \varphi_j(x) \\
& = \varphi'_{k+1} \left(i_{k+1} \varphi_{k+2}\right) \varphi_{k+3} ... \varphi_j(x) \\
& = \varphi'_{k+1} \left(\varphi'_{k+2} i_{k+2}\right) \varphi_{k+3} ... \varphi_j(x) \\
& = ... = \varphi'_{k+1} ... \varphi'_{j-1} \left(i_{j-1} \varphi_j\right)(x). \\
& = \varphi'_{k+1} ... \varphi'_{j-1} \left(\varphi'_j i_j\right)(x) = \left(\varphi'_{k+1} ... \varphi'_j\right) i_j(x) \\
& = E'_{k+1,\,j} i_j(x).
\end{aligned}$$

Therefore, the square of inclusions

$$\begin{array}{ccc}
A_k & \subset^{E_{k+1,\,j}} & A_j \\
\cup^{i_k} & & \cup^{i_j} \\
A'_k & \subset^{E'_{k+1,\,j}} & A'_j
\end{array}$$

is a commuting square of amalgamated NCPSpaces. ∎

Corollary 4.2. *Let*

$$\begin{array}{ccccccccccc}
\mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & A_1 & \subset^{\varphi_2} & A_2 & \subset^{\varphi_3} & A_3 & \subset^{\varphi_4} & \cdots \\
 & & \cup^{i_0} & & \cup^{i_1} & & \cup^{i_2} & & \cup^{i_3} & & \cdots \\
\mathbb{C} & \subset^{\varphi'_0} & B' & \subset^{\varphi'_1} & A'_1 & \subset^{\varphi'_2} & A'_2 & \subset^{\varphi'_3} & A'_3 & \subset^{\varphi'_4} & \cdots
\end{array}$$

be a commuting ladder of amalgamated NCPSpaces. Let $(j_k)_{k=1}^{\infty}$ *be any (sub)sequence of* $\mathbb{N}$. *Then we can construct a commuting (sub)ladder of amalgamated NCPSpaces*

(4.2.1)

$$\begin{array}{lllllllll}
\mathbb{C} & \subset^{E_{j_1}} & A_{j_1} & \subset^{E_{j_1+1,\,j_2}} & A_{j_2} & \subset^{E_{j_2+1,\,j_3}} & A_{j_3} & \subset & \cdots \\
 & & \cup^{i_{j_1}} & & \cup^{i_{j_2}} & & \cup^{i_{j_3}} & & \cdots \\
\mathbb{C} & \subset^{E'_{j_1}} & A'_{j_1} & \subset^{E'_{j_1+1,\,j_2}} & A'_{j_2} & \subset^{E'_{j_2+1,\,j_3}} & A'_{j_3} & \subset & \cdots.
\end{array}$$

Proof. By the previous proposition, we know that, for each chosen j_p ($p \in \mathbb{N}$),

$$\begin{array}{lll} A_{j_p} & \subset^{E_{j_p+1,\, j_{p+1}}} & A_{j_{p+1}} \\ \cup^{i_{j_p}} & & \cup^{i_{j_{p+1}}} \\ A'_{j_p} & \subset^{E'_{j_p+1,\, j_{p+1}}} & A'_{j_{p+1}} \end{array}$$

is a commuting square of amalgamated NCPSpaces. So the ladder (4.2.1) satisfies the condition (ii) of Definition 4.1. By the previous chapter (or by compatibility), we know that the given two towers are all towers of amalgamated NCPSpaces. So the condition (i) of Definition 4.1 is also satisfied. Thus the ladder (4.2.1) is a commuting ladder of amalgamated NCPSpaces. ∎

The above proposition and corollary show that if we have a **commuting** ladder of amalgamated NCPSpaces, then we can naturally define conditional expectations from the algebras, A_j's, in the upper tower of the ladder to the algebras, A'_k 's, in the lower tower of the ladder, whenever $k < j$. The commuting property is crucial. Hence we can observe the R-transform theory for those conditional expectations. Also, the above corollary shows us that if we have a commuting ladder of amalgamated NCPSpaces, then we can choose infinitely many commuting subladders of amalgamated NCPSpaces from the given ladder.

We observed, in the previous example, that we have the following commuting ladder of amalgamated (compressed) NCPSpaces ; Let (A, φ) be a NCPSpace over B and let D be a subalgebra of A containing B. Assume that we have a conditional expectation $E_D^A : A \rightarrow D$ defined by $E_B^A(a) = a$ if $a \in D$ and $E_D^A(a) = 0_D$, otherwise. Now, let $(p_k)_{k=1}^{\infty} \subset A_{pro}$ be a scalar-valued chain of projections such that $\varphi(p_k) = \alpha_k$, $\forall k \in \mathbb{N}$. Then we have the commuting ladder of amalgamated NCPSpaces,

(*)

$$\begin{array}{lllllllll} \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & p_1Ap_1 & \subset^{\varphi_2} & p_2Ap_2 & \subset^{\varphi_3} & \cdots \\ & & \cup^{E_D^A} & & \cup^{E_D^A} & & \cup^{E_D^A} & & \\ \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & p_1Dp_1 & \subset^{\varphi_2} & p_2Dp_2 & \subset^{\varphi_3} & \cdots . \end{array}$$

We will call this ladder, a commuting ladder of amalgamated compressed NCPSpaces induced by $(p_k)_{k=1}^{\infty}$ and E_D^A. Remark that it is possible that $D = p_NAp_N$, for some $N \in \mathbb{N}$. Then the above ladder can be regarded as

$$\begin{array}{lllllllll} \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & p_1Ap_1 & \subset^{\varphi_2} \cdots \subset^{\varphi_N} & p_NAp_N & \subset & \cdots \\ & & \cup^{E_D^A} & & \cup^{E_D^A} & & {}_{\shortparallel}{}^{E_D^A} & & \\ \mathbb{C} & \subset^{\varphi_0} & B & \subset^{\varphi_1} & p_1Dp_1 & \subset^{\varphi_2} \cdots \subset^{\varphi_N} & D & = & \cdots . \end{array}$$

i.e, we take $p_NDp_N = p_{N+1}Dp_{N+1} = p_{N+2}Dp_{N+2} = ... = D$. In this case, we have that

$$\varphi_{N+j} \mid_{p_{N+j}Dp_{N+j}} : p_{N+j}Dp_{N+j} \rightarrow p_{N+(j-1)}Dp_{N+(j-1)}$$

as the identity map, $i_D : D \to D$, for all $j \in \mathbb{N}$.

Proposition 4.3. *Let B be a unital algebra and (A, φ), a NCPSpace over B and let D be a subalgebra of A containing B. Suppose that we have a scaalr-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$, with $\varphi(p_k) = \alpha_k \cdot 1_B$, $\forall k$, and a conditional expectation $E_D^A : A \to D$ defined by $E_D^A(x) = x$, if $x \in D$ and $E_D^A(x) = 0_D = 0_B$, otherwise. Assume that we have a commuting ladder of amalgamated compressed NCPSpaces (*). Then, for any $k < j \in \mathbb{N}$, we can define a conditional expectation $F_{k+1,\,j} : p_j A p_j \to p_k D p_k$ satisfying the following property ; let X and Y be subsets of A. If $X \cup Y$ and $\{p_k\}$ are free over B, in (A, φ), then X and Y are free over $p_k A p_k$, in (A, φ) if and only if $p_j X p_j$ and $p_j Y p_j$ are free over $p_k D p_k$, in $\left(p_j A p_j,\, \frac{1}{\alpha_{k+1}} F_{k+1,\,j}\right)$.*

Proof. Suppose that we have a ladder of amalgamated compressed NCPSpaces (*) induced by $(p_k)_{k=1}^{\infty} \subset A_{pro}$ and a canonical conditional expectation, $E_D^A : A \to D$. Then we can define a conditional expectation,

$$F_{k+1,\,j} : p_j A p_j \to p_k D p_k,$$

for any $k < j$, by

$$F_{k+1,\,j} = E_D^A \left(\frac{1}{\alpha_{k+1}} \cdot E_{k+1,\,j} \right).$$

Recall that if X and Y are free over $p_k A p_k$, in (A, φ), then $p_j X p_j$ and $p_j Y p_j$ are free over $p_k A p_k$, in $\left(p_j A p_j,\, \frac{1}{\alpha_{k+1}} E_{k+1,\,j}\right)$ (See Section 2.3). Also, since

$$E_D^A E_{k+1,\,j} = E_{k+1,\,j}\, E_D^A,$$

we need to show that $p_j \left(E_D^A(X)\right) p_j$ and $p_j \left(E_D^A(Y)\right) p_j$ are free over $p_k D p_k$. But, trivially, if X and Y are free over $p_k A p_k$, then $E_D^A(X)$ and $E_D^A(Y)$ are free over $p_k D p_k$, by the very definition of E_D^A. i.e,

$$E_D^A(X) = \{x \in D : x \in D \cap X\} \text{ and } E_D^A(Y) = \{y \in D : y \in D \cap Y\}.$$

Thus they are automaticlly free under the hypothesis. So, we have that, if X and Y are free over $p_k A p_k$, in (A, φ) , then

(i) $p_j X p_j$ and $p_j Y p_j$ are free over $p_k A p_k$, in $\left(p_j A p_j,\, \frac{1}{\alpha_{k+1}} E_{k+1,\,j}\right)$

(ii) By (i), $p_j E_D^A(X) p_j$ and $p_j E_D^A(Y) p_j$ are also free over $p_k D p_k$ in $\left(p_j A p_j,\ \frac{1}{\alpha_{k+1}} E_D^A E_{k+1,\,j} \right)$.

So, we can conclude that $p_k X p_k$ and $p_k Y p_k$ are free over $p_k D p_k$, in $(p_j A p_j,\ F_{k+1,\,j})$. ∎

As an application, we have the following theorem ;

Theorem 4.4. *Let B be a unital algebra and (A, φ), a NCPSpace over B and let D be a subalgebra of A containing B. Suppose that we have a scaalr-valued chain of projections $(p_k)_{k=1}^{\infty} \subset A_{pro}$, with $\varphi(p_k) = \alpha_k \cdot 1_B$, $\forall k$, and a conditional expectation $E_D^A : A \to D$ defined by $E_D^A(x) = x$, if $x \in D$ and $E_D^A(x) = 0_D = 0_B$, otherwise. Assume that we have a commuting ladder of amalgamated compressed NCPSpaces (*). Let $F_{k+1,\,j} : p_j A p_j \to p_k A p_k$ be $E_D^A E_{k+1,\,j}$. Assume that we have two $p_k A p_k$-free subsets $X = \{p_j x_1 p_j, ..., p_j x_s p_j\}$ and $Y = \{p_j y_1 p_j, ..., p_j y_s p_j\}$. If $X \cup Y$ and $\{p_j\}$ is free over $p_k A p_k$, in $(p_j A p_j, E_{k+1,\,j})$, then, for any fixed $k < j \in \mathbb{N}$, we have that*

(1) $$R^{(F_{k+1,\,j})}_{p_j x_1 p_j, ..., p_j x_s p_j, p_j y_1 p_j, ..., p_j y_s p_j}(z_1, ..., z_{2s}) = R^{(F_{k+1,\,j})}_{p_j x_1 p_j, ..., p_j x_s p_j}(z_1, ..., z_s) + R^{(F_{k+1,\,j})}_{p_j y p_j, ..., p_j y_s p_j}(z_{s+1}, ..., z_{2s}).$$

(2) $$R^{(F_{k+1,\,j})}_{p_j x_1 p_j + p_j y_1 p_j, ..., p_j x_s p_j + p_j y_s p_j}(z_1, ..., z_{2s}) = R^{(F_{k+1,\,j})}_{p_j x_1 p_j, ..., p_j x_s p_j}(z_1, ..., z_s) + R^{(F_{k+1,\,j})}_{p_j y p_j, ..., p_j y_s p_j}(z_1, ..., z_s).$$

(3) $$\begin{aligned} R^{(F_{k+1,\,j})}_{(p_j x_1 p_j)(p_j y_1 p_j), ..., (p_j x_s p_j)(p_j y_s p_j)}(z_1, ..., z_s) &= \left(R^{(F_{k+1,\,j})}_{p_j x_1 p_j, ..., p_j x_s p_j} \ \boxed{*}_B \ R^{(F_{k+1,j})\,:\,t}_{p_j y_1 p_j, ..., p_j y_s p_j} \right)(z_1, ..., z_s) \\ &= \left(R^{(F_{k+1,\,j})}_{p_j x_1 p_j, ..., p_j x_s p_j} \ \boxed{*}_B \ R^{(\varphi)\,:\,t}_{y_1, \,.\,.\,.\,, y_s} \right)(z_1, ..., z_s). \end{aligned}$$

□

We can extend this construction of such ladders from two pairs of towers of amalgamated NCPSpaces to nets of amalgamated NCPSpaces such as

$$\begin{array}{ccccccccccc} \mathbb{C} & \subset & B & \subset & A_1 & \subset & A_2 & \subset & A_3 & \subset & \cdots \\ & & \cup & & \cup & & \cup & & \cup & & \\ \mathbb{C} & \subset & B' & \subset & A_1' & \subset & A_2' & \subset & A_3' & \subset & \cdots \\ & & \cup & & \cup & & \cup & & \cup & & \\ \mathbb{C} & \subset & B'' & \subset & A_1'' & \subset & A_2'' & \subset & A_3'' & \subset & \cdots \\ & & \cup & & \cup & & \cup & & \cup & & \\ & & \vdots & & \vdots & & \vdots & & \vdots & & \ddots \end{array}$$

But even in the commuting ladders, we have seen that it is difficult to consider the amalgamated freeness between two algebras in that ladderWe only considered the case when we have a commuting ladder of amalgamated compressed NCPSpaces induced by a scalar-valued chain of projections and the given "good" conditional expectation.

References

[1] A. Nica, R-transform in Free Probability, IHP course note, available at www.math.uwaterloo.ca/˜anica.

[2] A. Nica, R-transforms of Free Joint Distributions and Non-crossing Partitions, J. of Func. Anal, 135 (1996), 271-296.

[7] A. Nica, D. Shlyakhtenko and R. Speicher, R-cyclic Families of Matrices in Free Probability, J. of Funct Anal, 188 (2002), 227-271.

[4] A. Nica, D. Shlyakhtenko and R. Speicher, R-diagonal Elements and Freeness with Amalgamation, Canad. J. Math. Vol 53, Num 2, (2001) 355-381.

[5] D. Shlyakhtenko, Some Applications of Freeness with Amalgamation, J. Reine Angew. Math, 500 (1998), 191-212.

[6] D. Shlyakhtenko, A-Valued Semicircular Systems, J. of Funct Anal, 166 (1999), 1-47.

[7] D. Voiculescu, Operations on Certain Non-commuting Operator-Valued Random Variables, Astérisque, 232 (1995), 243-275.

[8] D.Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).

[9] I. Cho, Amalgamated Boxed Convolution and Amalgamated R-transform Theory (preprint).

[10] I. Cho, Compressed Amalgamated R-transform Theory, preprint.

[11] I. Cho, Perturbed R-transform Theory, preprint.

[12] I. Cho, Compatibility of a noncommutative probability space and an amalgamated noncommutative probability space, preprint

[13] R. Speicher, Combinatorics of Free Probability Theory IHP course note, available at www.mast.queensu.ca/˜speicher.

[14] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132 , Num 627 , (1998).

Dep. of Math, Univ of Iowa, Iowa Sity, IA, U.S.A
E-mail address: `icho@math.uiowa.edu`